Random Convex Hull and Extreme Value Statistics

Satya N. Majumdar

Laboratoire de Physique Théorique et Modèles Statistiques,CNRS,
Université Paris-Sud, France

Collaborators:

A. Comtet (LPTMS, Orsay, FRANCE)
J. Randon-Furling (Univ. Paris-1, FRANCE)

Ref: Phys. Rev. Lett. 103, 140602 (2009)

Extended Review: J. Stat. Phys. 138, 955 (2010)

Plan

Plan:

- Random Convex Hull $\Longrightarrow$ definition

Plan

Plan:

- Random Convex Hull $\Longrightarrow$ definition
- Convex Hull of n planar Brownian motions

Plan

Plan:

- Random Convex Hull $\Longrightarrow$ definition
- Convex Hull of n planar Brownian motions
- Motivation $\Longrightarrow$ an ecological problem

Plan

Plan:

- Random Convex Hull $\Longrightarrow$ definition
- Convex Hull of n planar Brownian motions
- Motivation $\Longrightarrow$ an ecological problem
- Cauchy's formulae for perimeter and area of a closed convex curve in two dimensions

Plan

Plan:

- Random Convex Hull $\Longrightarrow$ definition
- Convex Hull of n planar Brownian motions
- Motivation $\Longrightarrow$ an ecological problem
- Cauchy's formulae for perimeter and area of a closed convex curve in two dimensions

  $\Longrightarrow$ applied to random convex polygon

Plan

Plan:

- Random Convex Hull $\Longrightarrow$ definition
- Convex Hull of n planar Brownian motions
- Motivation $\Longrightarrow$ an ecological problem
- Cauchy's formulae for perimeter and area of a closed convex curve in two dimensions

  $\Longrightarrow$ applied to random convex polygon

  $\Longrightarrow$ link to Extreme Value Statistics

Plan

Plan:

- Random Convex Hull $\Longrightarrow$ definition
- Convex Hull of n planar Brownian motions
- Motivation $\Longrightarrow$ an ecological problem
- Cauchy's formulae for perimeter and area of a closed convex curve in two dimensions

  $\Longrightarrow$ applied to random convex polygon

  $\Longrightarrow$ link to Extreme Value Statistics
- Exact results for the mean perimeter and the mean area for all n.

Plan

Plan:

- Random Convex Hull $\Longrightarrow$ definition
- Convex Hull of n planar Brownian motions
- Motivation $\Longrightarrow$ an ecological problem
- Cauchy's formulae for perimeter and area of a closed convex curve in two dimensions

  $\Longrightarrow$ applied to random convex polygon

  $\Longrightarrow$ link to Extreme Value Statistics
- Exact results for the mean perimeter and the mean area for all n.
- Summary and Conclusions

Shape of a set of Points





Shape of a set of Points: Convex Hull



Random Convex Hull in a Plane





- Convex Hull $\Longrightarrow$ Minimal convex polygon enclosing the set

Random Convex Hull in a Plane





- Convex Hull $\Longrightarrow$ Minimal convex polygon enclosing the set
- The shape of the convex hull $\rightarrow$ different for each sample

Random Convex Hull in a Plane





- Convex Hull $\Longrightarrow$ Minimal convex polygon enclosing the set
- The shape of the convex hull $\rightarrow$ different for each sample
- Points drawn from a distribution $\rightarrow$ Independent or Correlated

Random Convex Hull in a Plane





- Convex Hull $\Longrightarrow$ Minimal convex polygon enclosing the set
- The shape of the convex hull $\rightarrow$ different for each sample
- Points drawn from a distribution $\rightarrow$ Independent or Correlated
- Question: Statistics of observables: perimeter, area and no. of vertices

Independent Points in a Plane



Each point chosen independently from the same distribution

Independent Points in a Plane



Each point chosen independently from the same distribution

Associated Random Convex Hull $\rightarrow$ well studied by diverse methods

Independent Points in a Plane



Each point chosen independently from the same distribution

Associated Random Convex Hull $\rightarrow$ well studied by diverse methods

P. Lévy ('48), J. Geffroy ('59), Spitzer & Widom ('59), Baxter ('59)..
Rényi & Sulanke ('63), Efron ('65), ...many others

Correlated Points: Vertices of an Open Random Walk



8–step walk



another 8–step walk

- Continuous-time limit: Brownian path of duration T

Correlated Points: Vertices of an Open Random Walk



8–step walk



another 8–step walk

- Continuous-time limit: Brownian path of duration T
- mean perimeter and mean area of the associated Convex hull?

Correlated Points: Vertices of an Open Random Walk



8–step walk



another 8–step walk

- Continuous-time limit: Brownian path of duration T
- mean perimeter and mean area of the associated Convex hull?
- mean perimeter: $\langle L_1 \rangle = \sqrt{8\pi T}$ (Takács, '80)
- mean area: $\langle A_1 \rangle = \frac{\pi}{2} T$ (El Bachir, '83, Letac '93)

Correlated Points: Vertices of a Closed Random Walk



8 step random bridge



another 8 step bridge

• Continuous-time limit: Brownian bridge of duration T : starting at O and returning to it after time T

Correlated Points: Vertices of a Closed Random Walk



8 step random bridge



another 8 step bridge

- Continuous-time limit: Brownian bridge of duration T : starting at O and returning to it after time T
- mean perimeter: $\langle L_1 \rangle = \sqrt{\frac{\pi^3}{2}\, T}$ (Goldman, '96).

Correlated Points: Vertices of a Closed Random Walk



8 step random bridge



another 8 step bridge

- Continuous-time limit: Brownian bridge of duration T : starting at O and returning to it after time T
- mean perimeter: $\langle L_1 \rangle = \sqrt{\frac{\pi^3}{2}\, T}$ (Goldman, '96).
- mean area: $\langle A_1 \rangle =$?

Home Range Estimate via Convex Hull



Home Range Estimate via Convex Hull



Models of home range for animal movement, Worton (1987)
Integrating Scientific Methods with Habitat Conservation Planning, Murphy and Noon (1992)
Theory of home range estimation from displacement measurements of animal populations, Giuggioli et. al. (2005)
Home Range Estimates, Boyle et. al., (2009)

Home Range Estimate via Convex Hull



Models of home range for animal movement, Worton (1987)
Integrating Scientific Methods with Habitat Conservation Planning, Murphy and Noon (1992)
Theory of home range estimation from displacement measurements of animal populations, Giuggioli et. al. (2005)
Home Range Estimates, Boyle et. al., (2009)

Global Convex Hull of n Independent Brownian Paths





Global Convex Hull of n Independent Brownian Paths





• Mean perimeter $\langle L_n \rangle$ and mean area $\langle A_n \rangle$ of n independent Brownian paths (bridges) each of duration T?

Global Convex Hull of n Independent Brownian Paths





- Mean perimeter $\langle L_n \rangle$ and mean area $\langle A_n \rangle$ of n independent Brownian paths (bridges) each of duration T?
- $\langle L_n \rangle = \alpha_n \sqrt{T}$; $\langle A_n \rangle = \beta_n T$

Global Convex Hull of n Independent Brownian Paths





- Mean perimeter $\langle L_n \rangle$ and mean area $\langle A_n \rangle$ of n independent Brownian paths (bridges) each of duration T?
- $\langle L_n \rangle = \alpha_n \sqrt{T}$; $\langle A_n \rangle = \beta_n T$
- Recall $\boxed{\alpha_1 = \sqrt{8\pi}, \ \beta_1 = \pi/2}$ (open path)

$\boxed{\alpha_1 = \sqrt{\pi^3/2}, \ \beta_1 = ?}$ (closed path)

Global Convex Hull of n Independent Brownian Paths





- Mean perimeter $\langle L_n \rangle$ and mean area $\langle A_n \rangle$ of n independent Brownian paths (bridges) each of duration T?
- $\langle L_n \rangle = \alpha_n \sqrt{T}$; $\langle A_n \rangle = \beta_n T$
- Recall $\boxed{\alpha_1 = \sqrt{8\pi},\ \beta_1 = \pi/2}$ (open path)

  $\boxed{\alpha_1 = \sqrt{\pi^3/2},\ \beta_1 = ?}$ (closed path)
- $\boxed{\alpha_n,\ \beta_n = ?}$ $\rightarrow$ both for open and closed paths $\rightarrow$ n-dependence?

Global Convex Hull of n Independent Brownian Paths



$n = 3$ closed paths



$n = 10$ open paths

Cauchy's Formulae for a Closed Convex Curve



C : CLOSED CONVEX CURVE

- For any point $[X(s), Y(s)]$ on C define:

  Support function: $M(\theta) = \max_{s \in C} [X(s)\cos(\theta) + Y(s)\sin(\theta)]$

Cauchy's Formulae for a Closed Convex Curve



C : CLOSED CONVEX CURVE

- For any point $[X(s), Y(s)]$ on C define:

  Support function: $M(\theta) = \max_{s \in C} [X(s)\cos(\theta) + Y(s)\sin(\theta)]$

- Perimeter: $$\boxed{L = \int_0^{2\pi} d\theta \; M(\theta)}$$

- Area: $$\boxed{A = \frac{1}{2}\int_0^{2\pi} d\theta \; \left[M^2(\theta) - [M'(\theta)]^2\right]}$$

A simple physicist's proof of Cauchy's formula





A simple physicist's proof of Cauchy's formula



approximate C

by a polygon



$$M(\theta) = R\cos\theta$$

A simple physicist's proof of Cauchy's formula



approximate C

by a polygon



$$M(\theta) = R\cos\theta$$

Perimeter: $\int_{-\phi_1}^{\phi_2} M(\theta)d\theta = R\left[\sin(\phi_1) + \sin(\phi_2)\right] = L_{P_1AP_2}$

A simple physicist's proof of Cauchy's formula





$$M(\theta) = R\cos\theta$$

Perimeter: $\int_{-\phi_1}^{\phi_2} M(\theta)d\theta = R\left[\sin(\phi_1) + \sin(\phi_2)\right] = L_{P_1AP_2}$

Area: $\frac{1}{2}\int_{-\phi_1}^{\phi_2}\left[M^2(\theta) - (M'(\theta))^2\right]d\theta$

A simple physicist's proof of Cauchy's formula





$$\boxed{M(\theta) = R\cos\theta}$$

Perimeter: $\int_{-\phi_1}^{\phi_2} M(\theta)d\theta = R\left[\sin(\phi_1) + \sin(\phi_2)\right] = L_{P_1AP_2}$

Area: $\frac{1}{2}\int_{-\phi_1}^{\phi_2}\left[M^2(\theta) - (M'(\theta))^2\right]d\theta$

$= \frac{R^2}{2}\left[\sin(\phi_2)\cos(\phi_2) + \sin(\phi_1)\cos(\phi_1)\right] = A_{OP_1AP_2}$

Examples

a circle centered at the origin:

$$M(\theta) = r$$



Examples

a circle centered at the origin:

$$M(\theta) = r$$



$$L = \int_0^{2\pi} d\theta \; M(\theta) = 2\pi r$$

$$A = \frac{1}{2}\int_0^{2\pi} d\theta \; \left[M^2(\theta) - [M'(\theta)]^2\right] = \pi r^2$$

Examples

a circle centered at the origin:

$$M(\theta) = r$$



a circle touching the origin:

$$M(\theta) = r(1 + \sin\theta)$$



Examples

a circle centered at the origin:

$$M(\theta) = r$$



a circle touching the origin:

$$M(\theta) = r(1 + \sin\theta)$$



$$L = \int_0^{2\pi} d\theta \; M(\theta) = 2\pi r$$

$$A = \frac{1}{2}\int_0^{2\pi} d\theta \; \left[M^2(\theta) - [M'(\theta)]^2\right] = \pi r^2$$

Cauchy's formulae Applied to Convex Polygon



Let $(x_k, y_k) \in I \Longrightarrow$ vertices of an N-step random walk starting at O

Let C (green) be the associated Convex Hull

Cauchy's formulae Applied to Convex Polygon



$(x_k, y_k) \in I \Longrightarrow$ vertices of the walk

$C \rightarrow$ Convex Hull with coordinates $\{X(s), Y(s)\}$ on C

Cauchy's formulae Applied to Convex Polygon



$$M(\theta) = \max_{s \in C} [X(s)\cos\theta + Y(s)\sin\theta]$$

$$= \max_{k \in I} [x_k \cos\theta + y_k \sin\theta]$$

$$= x_{k^*} \cos\theta + y_{k^*} \sin\theta$$

$k^* \rightarrow$ label of the point with largest projection along θ

Support Function of a Convex Hull



$$M(\theta) = x_{k^*} \cos\theta + y_{k^*} \sin\theta$$
$$M'(\theta) = -x_{k^*} \sin\theta + y_{k^*} \cos\theta$$

Cauchy's Formulae Applied to Random Convex Hull



Cauchy's Formulae Applied to Random Convex Hull



Mean perimeter of a random convex polygon

$$\langle L \rangle = \int_0^{2\pi} d\theta \, \langle M(\theta) \rangle$$

with $M(\theta) = x_{k^*} \cos\theta + y_{k^*} \sin\theta$

Mean area of a random convex polygon

$$\langle A \rangle = \frac{1}{2} \int_0^{2\pi} d\theta \left[\langle M^2(\theta) \rangle - \langle [M'(\theta)]^2 \rangle \right]$$

with $M'(\theta) = -x_{k^*} \sin\theta + y_{k^*} \cos\theta$

Isotropically Distributed Points



Mean Perimeter

$$\langle L \rangle = 2\pi \langle M(0) \rangle$$

with $M(\theta = 0) = \max_{k \in I} \{x_k\} = x_{k*}$

Isotropically Distributed Points



Mean Perimeter

$$\langle L \rangle = 2\pi \langle M(0) \rangle$$

with $M(\theta = 0) = \max_{k \in I} \{x_k\} = x_{k^*}$

Mean Area

$$\langle A \rangle = \pi \left[\langle M^2(0) \rangle - \langle [M'(0)]^2 \rangle \right]$$

with $M'(\theta = 0) = y_{k^*}$

Isotropically Distributed Points



Mean Perimeter

$$\langle L \rangle = 2\pi \langle M(0) \rangle$$

with $M(\theta = 0) = \max_{k \in I} \{x_k\} = x_{k^*}$

Mean Area

$$\langle A \rangle = \pi \left[\langle M^2(0) \rangle - \langle [M'(0)]^2 \rangle \right]$$

with $M'(\theta = 0) = y_{k^*}$

$\Longrightarrow$ Link to Extreme Value Statistics

Cauchy's Formulae Applied to the Convex Hull of a Brownian Path (n=1)



$x(\tau)$, $y(\tau) \rightarrow$ a pair of independent one-dimensional Brownian motions: $0 \leq \tau \leq T$

$\frac{dx}{d\tau} = \eta_x(\tau)$
$\frac{dy}{d\tau} = \eta_y(\tau)$

Mean Perimeter

$$\langle L \rangle = 2\pi \langle M(0) \rangle$$

with $M(0) = \max_{0 \leq \tau \leq T} \{x(\tau)\} \equiv x(\tau^*)$

Mean Area

$$\langle A \rangle = \pi \left[\langle M^2(0) \rangle - \langle [M'(0)]^2 \rangle \right]$$

with $M'(0) = y(\tau^*)$

$M'(0) \to$ value of y at the special time τ^* when $x(\tau)$ is maximal





$M'(0) \to$ value of y at the special time τ^* when $x(\tau)$ is maximal





- $\langle M(0) \rangle = \int_0^\infty dM\ M\ \sigma_1(M|T)$; $\langle M^2(0) \rangle = \int_0^\infty dM\ M^2\ \sigma_1(M|T)$

$M'(0) \to$ value of y at the special time τ^* when $x(\tau)$ is maximal



- $\langle M(0) \rangle = \int_0^\infty dM\ M\ \sigma_1(M|T)$; $\langle M^2(0) \rangle = \int_0^\infty dM\ M^2\ \sigma_1(M|T)$
- $\sigma_1(M|T) \to$ prob. density of maximum $M(0)$ of $x(\tau)$ in $[0, T]$

$M'(0) \to$ value of y at the special time τ^* when $x(\tau)$ is maximal



- $\langle M(0) \rangle = \int_0^\infty dM\ M\ \sigma_1(M|T)$; $\langle M^2(0) \rangle = \int_0^\infty dM\ M^2\ \sigma_1(M|T)$
- $\sigma_1(M|T) \to$ prob. density of maximum $M(0)$ of $x(\tau)$ in $[0, T]$
- $\langle [M'(0)]^2 \rangle = \int_0^T d\tau^*\ \rho_1(\tau^*|T)\ \langle y^2(\tau^*) \rangle$

$M'(0) \to$ value of y at the special time τ^* when $x(\tau)$ is maximal



- $\langle M(0) \rangle = \int_0^\infty dM\ M\ \sigma_1(M|T)$; $\langle M^2(0) \rangle = \int_0^\infty dM\ M^2\ \sigma_1(M|T)$
- $\sigma_1(M|T) \to$ prob. density of maximum $M(0)$ of $x(\tau)$ in $[0, T]$
- $\langle [M'(0)]^2 \rangle = \int_0^T d\tau^*\ \rho_1(\tau^*|T)\ \langle y^2(\tau^*) \rangle = \langle \tau^* \rangle$

Distribution of M and τ^* for a single Brownian Path



Distribution of M and τ^* for a single Brownian Path



- Joint Distribution: $P_1(M, \tau^* | T) = \frac{M}{\pi \tau^{*3/2} \sqrt{T - \tau^*}} e^{-M^2/2\tau^*}$

Distribution of M and τ^* for a single Brownian Path



- Joint Distribution: $P_1(M, \tau^* | T) = \frac{M}{\pi \tau^{*3/2} \sqrt{T - \tau^*}} e^{-M^2/2\tau^*}$
- Marginals: $\sigma_1(M|T) = \sqrt{\frac{2}{\pi T}} e^{-M^2/2T}$

Distribution of M and τ^* for a single Brownian Path



- Joint Distribution: $P_1(M, \tau^* | T) = \frac{M}{\pi \tau^{*3/2} \sqrt{T-\tau^*}} e^{-M^2/2\tau^*}$
- Marginals: $\sigma_1(M|T) = \sqrt{\frac{2}{\pi T}} e^{-M^2/2T}$

$$\rho_1(\tau^*|T) = \frac{1}{\pi \sqrt{\tau^*(T-\tau^*)}} \rightarrow \text{Lévy's arcsine law}$$

Distribution of the time τ^* at which a Brownian Motion is maximal over [0,T]

Lévy's Arcsine Law: $\rho_1(\tau^*|T) = \frac{1}{T}\, f_1\left(\frac{\tau^*}{T}\right)$

$$f_1(x) = \frac{1}{\pi\sqrt{x(1-x)}}$$

Distribution of the time τ^* at which a Brownian Motion is maximal over [0,T]

Lévy's Arcsine Law: $\rho_1(\tau^*|T) = \frac{1}{T}\, f_1\left(\frac{\tau^*}{T}\right)$

$$f_1(x) = \frac{1}{\pi\sqrt{x(1-x)}}$$

Cumulative distribution: $\mathrm{Prob}(\tau^* \leq t|T) = \frac{2}{\pi} \arcsin\left(\sqrt{t}\right)$



Results for n=1 Open Brownian Path



$x(\tau)$, $y(\tau) \rightarrow$ a pair of independent one-dimensional Brownian motions over $0 \leq \tau \leq T$

Mean Perimeter

$$\langle L \rangle = \sqrt{8\pi T}$$

Mean Area

$$\langle A \rangle = \frac{\pi T}{2}$$

Takács, *Expected perimeter length*, Amer. Math. Month., **87** (1980)
El Bachir, (1983)

Results for n=1 Closed Brownian Path



$x(\tau)$, $y(\tau) \rightarrow$ a pair of independent one-dimensional Brownian bridges over $0 \leq \tau \leq T$

Mean Perimeter

$\langle L \rangle = \sqrt{\frac{\pi^3 T}{2}}$ Goldman, '96

Results for n=1 Closed Brownian Path



$x(\tau)$, $y(\tau) \rightarrow$ a pair of independent one-dimensional Brownian bridges over $0 \leq \tau \leq T$

Mean Perimeter

$\langle L \rangle = \sqrt{\frac{\pi^3 T}{2}}$ Goldman, '96

Mean Area

$\boxed{\langle A \rangle = \frac{\pi T}{3}}$ $\rightarrow$ New Result

Convex Hull of n Independent Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$ independent one-dimensional Brownian paths each of duration T

Mean Perimeter

$$\langle L_n \rangle = 2\pi \langle M_n \rangle$$

with $M_n = \max_{\tau,i} \{x_i(\tau)\} \equiv x_{i*}(\tau^*)$

Mean Area

$$\langle A_n \rangle = \pi \left[\langle M_n^2 \rangle - \langle [M_n']^2 \rangle \right]$$

with $M_n' = y_{i*}(\tau^*)$

Distribution of the global maximum M and τ^* for n paths



Distribution of the global maximum M and τ^* for n paths



- Joint Distribution: $P_n(M, \tau^* | T) = n\, P_1(M, \tau^* | T) \left[\mathrm{erf}\left(\frac{M}{\sqrt{2T}}\right)\right]^{n-1}$

$$\mathrm{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z du\ e^{-u^2}$$

Marginals of M and τ^* for arbitrary n

- Marginals: $\sigma_n(M|T) = \sqrt{\frac{2}{\pi T}}\, n\, e^{-M^2/2T} \left[\mathrm{erf}\left(\frac{M}{\sqrt{2T}}\right)\right]^{n-1}$

Marginals of M and τ^* for arbitrary n

- Marginals: $\sigma_n(M|T) = \sqrt{\frac{2}{\pi T}}\, n\, e^{-M^2/2T} \left[\mathrm{erf}\left(\frac{M}{\sqrt{2T}}\right)\right]^{n-1}$

$$\rho_n(\tau^*|T) = \frac{1}{T} f_n\left(\tau^*/T\right)$$

Marginals of M and τ^* for arbitrary n

- Marginals: $\sigma_n(M|T) = \sqrt{\frac{2}{\pi T}}\, n\, e^{-M^2/2T} \left[\mathrm{erf}\left(\frac{M}{\sqrt{2T}}\right)\right]^{n-1}$

$$\rho_n(\tau^*|T) = \frac{1}{T} f_n\left(\tau^*/T\right)$$

$$f_n(x) = \frac{2n}{\pi\sqrt{x(1-x)}} \int_0^\infty u\, e^{-u^2} \left[\mathrm{erf}(u\sqrt{x})\right]^{n-1} du$$

Marginals of M and τ^* for arbitrary n

- Marginals: $\sigma_n(M|T) = \sqrt{\frac{2}{\pi T}}\, n\, e^{-M^2/2T} \left[\mathrm{erf}\left(\frac{M}{\sqrt{2T}}\right)\right]^{n-1}$

$$\rho_n(\tau^*|T) = \frac{1}{T} f_n\left(\tau^*/T\right)$$

$$f_n(x) = \frac{2n}{\pi\sqrt{x(1-x)}} \int_0^\infty u\, e^{-u^2} \left[\mathrm{erf}(u\sqrt{x})\right]^{n-1} du$$



Marginals for n Independent Brownian Bridges





Marginals for n Independent Brownian Bridges





- Marginals: $\sigma_n(M|T) = \frac{4n}{T}\, M \left(1 - e^{-2M^2/T}\right)^{n-1}$

Marginals for n Independent Brownian Bridges





- Marginals: $\sigma_n(M|T) = \frac{4n}{T}\, M \left(1 - e^{-2M^2/T}\right)^{n-1}$

$$\rho_n(\tau^*|T) = \frac{1}{T}\, g_n\left(\tau^*/T\right)$$

Marginals for n Independent Brownian Bridges





- Marginals: $\sigma_n(M|T) = \frac{4n}{T}\, M \left(1 - e^{-2M^2/T}\right)^{n-1}$

$$\rho_n(\tau^*|T) = \frac{1}{T}\, g_n\left(\tau^*/T\right)$$

$$g_n(x) = n \sum_{k=0}^{n-1} \binom{n-1}{k} \frac{(-1)^k}{[1 + 4\,k\,x\,(1-x)]^{\frac{3}{2}}}$$

Mean Perimeter and Mean Area of the Convex Hull of n Independent Open Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$ independent one-dimensional Brownian paths over $0 \leq \tau \leq T$

Mean Perimeter (open paths)

$$\langle L_n \rangle = 2\pi \langle M_n \rangle$$

with $M_n = \max_{\tau,i} \{x_i(\tau)\} \equiv x_{i*}(\tau^*)$

Mean Perimeter and Mean Area of the Convex Hull of n Independent Open Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$ independent one-dimensional Brownian paths over $0 \leq \tau \leq T$

Mean Perimeter (open paths)

$$\langle L_n \rangle = \alpha_n \sqrt{T}$$

$$\alpha_n = 4n\sqrt{2\pi} \int_0^\infty du\ u\ e^{-u^2}\ [\mathrm{erf}(u)]^{n-1}$$

Mean Perimeter and Mean Area of the Convex Hull of n Independent Open Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$
independent
one-dimensional Brownian
paths over $0 \leq \tau \leq T$

Mean Perimeter (open paths)

$$\langle L_n \rangle = \alpha_n \sqrt{T}$$

$$\boxed{\alpha_n = 4n\sqrt{2\pi} \int_0^\infty du\ u\ e^{-u^2}\ [\mathrm{erf}(u)]^{n-1}}$$

$$\alpha_1 = \sqrt{8\pi} = 5,013..$$

$$\alpha_2 = 4\sqrt{\pi} = 7,089..$$

$$\alpha_3 = 24\,\frac{\tan^{-1}\left(1/\sqrt{2}\right)}{\sqrt{\pi}} = 8,333..$$

Mean Perimeter and Mean Area of the Convex Hull of n Independent Open Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$
independent
one-dimensional Brownian
paths over $0 \leq \tau \leq T$

Mean Area (open paths)

$$\langle A_n \rangle = \pi \left[\langle M_n^2 \rangle - \langle [M_n']^2 \rangle \right]$$

with $M_n' = y_{i*}(\tau^*)$

Mean Perimeter and Mean Area of the Convex Hull of n Independent Open Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$ independent one-dimensional Brownian paths over $0 \le \tau \le T$

Mean Area (open paths)

$$\langle A_n \rangle = \beta_n T$$

$$\beta_n = 4n\sqrt{\pi} \int_0^\infty du\; u\; [\mathrm{erf}(u)]^{n-1} \left(u e^{-u^2} - h(u) \right)$$

$$h(u) = \frac{1}{2\sqrt{\pi}} \int_0^1 \frac{e^{-u^2/t}\, dt}{\sqrt{t(1-t)}}$$

Mean Perimeter and Mean Area of the Convex Hull of n Independent Open Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$
independent
one-dimensional Brownian
paths over $0 \leq \tau \leq T$

Mean Area (open paths)

$$\langle A_n \rangle = \beta_n T$$

$$\beta_n = 4n\sqrt{\pi} \int_0^\infty du\; u\; [\mathrm{erf}(u)]^{n-1} \left(u e^{-u^2} - h(u) \right)$$

$$h(u) = \frac{1}{2\sqrt{\pi}} \int_0^1 \frac{e^{-u^2/t}\, dt}{\sqrt{t(1-t)}}$$

$$\begin{aligned} \beta_1 &= \frac{\pi}{2} = 1,570.. \\ \beta_2 &= \pi = 3,141.. \\ \beta_3 &= \pi + 3 - \sqrt{3} = 4,409.. \end{aligned}$$

Mean Perimeter and Mean Area of the Convex Hull of n Independent Closed Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$ independent one-dimensional Brownian bridges over $0 \leq \tau \leq T$

Mean Perimeter (Closed Paths)

$$\langle L_n^c \rangle = \alpha_n^c \sqrt{T}$$

$$\alpha_n^c = \frac{\pi^{3/2}}{\sqrt{2}} \sum_{k=1}^{n} \binom{n}{k} \frac{(-1)^{k+1}}{\sqrt{k}}$$

Mean Perimeter and Mean Area of the Convex Hull of n Independent Closed Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$ independent one-dimensional Brownian bridges over $0 \leq \tau \leq T$

Mean Perimeter (Closed Paths)

$$\langle L_n^c \rangle = \alpha_n^c \sqrt{T}$$

$$\boxed{\alpha_n^c = \frac{\pi^{3/2}}{\sqrt{2}} \sum_{k=1}^{n} \binom{n}{k} \frac{(-1)^{k+1}}{\sqrt{k}}}$$

$$\begin{aligned} \alpha_1^c &= \sqrt{\pi^3/2} = 3,937. \\ \alpha_2^c &= \sqrt{\pi^3}(\sqrt{2} - 1/2) = 5,090.. \\ \alpha_3^c &= \sqrt{\pi^3}\left(\frac{3}{\sqrt{2}} - \frac{3}{2} + \frac{1}{\sqrt{6}}\right) = 5,732.. \end{aligned}$$

Mean Perimeter and Mean Area of the Convex Hull of n Independent Closed Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$ independent one-dimensional Brownian bridges over $0 \leq \tau \leq T$

Mean Area (Closed Paths)

$$\langle A_n^c \rangle = \beta_n^c\, T$$

$$\beta_n^c = \frac{\pi}{2}\left[\sum_{k=1}^{n} \frac{1}{k} - \frac{n}{3} + \frac{1}{2}\sum_{k=2}^{n} (-1)^k\, w(k)\right]$$

$$w(k) = \binom{n}{k}\,(k-1)^{-3/2}\left(k \tan^{-1}(\sqrt{k-1}) - \sqrt{k-1}\right)$$

Mean Perimeter and Mean Area of the Convex Hull of n Independent Closed Brownian Paths



$x_i(\tau)$, $y_i(\tau) \to 2\,n$ independent one-dimensional Brownian bridges over $0 \le \tau \le T$

Mean Area (Closed Paths)

$$\langle A_n^c \rangle = \beta_n^c\, T$$

$$\boxed{\beta_n^c = \frac{\pi}{2}\left[\sum_{k=1}^{n}\frac{1}{k} - \frac{n}{3} + \frac{1}{2}\sum_{k=2}^{n}(-1)^k\, w(k)\right]}$$

$$w(k) = \binom{n}{k}\,(k-1)^{-3/2}\left(k\tan^{-1}(\sqrt{k-1}) - \sqrt{k-1}\right)$$

$$\begin{aligned}\beta_1^c &= \frac{\pi}{3} = 1,047..\\ \beta_2^c &= \frac{\pi(4+3\pi)}{24} = 1,757..\\ \beta_3^c &= 2,250..\end{aligned}$$

Numerical Check



The coefficients α_n (mean perimeter) (lower triangle), β_n (mean area) (upper triangle) of n open paths and similarly α_n^c (square) and β_n^c (diamond) for n closed paths, plotted against n. The symbols denote numerical simulations (up to $n = 10$, with 10^3 realisations for each point)

Asymptotics for large n

For n open paths:

$$\langle L_n \rangle \simeq \left(2\,\pi\sqrt{2\ln n}\right)\ \sqrt{T}$$

$$\langle A_n \rangle \simeq (2\,\pi \ln n)\ T$$

Asymptotics for large n

For n open paths:

$$\langle L_n \rangle \simeq \left(2\,\pi\sqrt{2\ln n}\right)\ \sqrt{T}$$

$$\langle A_n \rangle \simeq (2\,\pi \ln n)\ \ T$$

For n closed paths:

$$\langle L_n^c \rangle \simeq \left(\pi\,\sqrt{2\ln n}\right)\ \sqrt{T}$$

$$\langle A_n^c \rangle \simeq \left(\tfrac{\pi}{2}\ln n\right)\ \ T$$

Asymptotics for large n

For n open paths:

$$\langle L_n \rangle \simeq \left(2\,\pi\sqrt{2\ln n}\right)\ \sqrt{T}$$

$$\langle A_n \rangle \simeq (2\,\pi \ln n)\ \ T$$

For n closed paths:

$$\langle L_n^c \rangle \simeq \left(\pi\,\sqrt{2\ln n}\right)\ \sqrt{T}$$

$$\langle A_n^c \rangle \simeq \left(\tfrac{\pi}{2}\ln n\right)\ \ T$$

- As $n \to \infty$, Convex Hull $\to$ Circle

Asymptotics for large n

For n open paths:

$$\langle L_n \rangle \simeq \left(2\,\pi\sqrt{2\ln n}\right)\ \sqrt{T}$$

$$\langle A_n \rangle \simeq (2\,\pi \ln n)\ \ T$$

For n closed paths:

$$\langle L_n^c \rangle \simeq \left(\pi\,\sqrt{2\ln n}\right)\ \sqrt{T}$$

$$\langle A_n^c \rangle \simeq \left(\tfrac{\pi}{2}\ln n\right)\ \ T$$

- As $n \to \infty$, Convex Hull $\to$ Circle
- Very slow growth with $n \Longrightarrow$ good news for conservation

Summary and Conclusion

- Unified approach adapting Cauchy's formulae

$\Longrightarrow$ Mean Perimeter and Area of Random Convex Hull

both for Independent and Correlated points

Summary and Conclusion

- Unified approach adapting Cauchy's formulae

  $\Longrightarrow$ Mean Perimeter and Area of Random Convex Hull

  both for Independent and Correlated points

- Provides a link Random Convex Hull $\Longrightarrow$ Extreme Value Statistics

Summary and Conclusion

- Unified approach adapting Cauchy's formulae

  $\Longrightarrow$ Mean Perimeter and Area of Random Convex Hull

  both for Independent and Correlated points

- Provides a link Random Convex Hull $\Longrightarrow$ Extreme Value Statistics

- Exact results for n planar Brownian paths $\rightarrow$ Open and Closed

Summary and Conclusion

- Unified approach adapting Cauchy's formulae

  $\Longrightarrow$ Mean Perimeter and Area of Random Convex Hull

  both for Independent and Correlated points

- Provides a link Random Convex Hull $\Longrightarrow$ Extreme Value Statistics

- Exact results for n planar Brownian paths $\rightarrow$ Open and Closed

  $\Longrightarrow$ Ecological Implication: Home Range Estimate

Very slow (logarithmic) growth of Home Range with population size n

Open Questions

- For n planar random walks each of N steps

  Mean no. of Vertices $\langle V_n(N) \rangle \to$?

Open Questions

- For n planar random walks each of N steps

  Mean no. of Vertices $\langle V_n(N) \rangle \to$?

  Only $n = 1$ case (Open) walk, the result is known:

$$\boxed{\langle V_1(N) \rangle \simeq 2 \left[1 + \frac{1}{2} + \frac{1}{3} + \ldots + \frac{1}{N} \right] \sim 2 \log(N)} \quad \text{for large } N$$

(Baxter, '61)

Open Questions

- For n planar random walks each of N steps

  Mean no. of Vertices $\langle V_n(N) \rangle \to$?

  Only $n = 1$ case (Open) walk, the result is known:

$$\boxed{\langle V_1(N) \rangle \simeq 2 \left[1 + \frac{1}{2} + \frac{1}{3} + \ldots + \frac{1}{N} \right] \sim 2 \log(N)} \quad \text{for large } N$$

(Baxter, '61)

- Distributions of the perimeter, area of the convex hull?

Open Questions

- For n planar random walks each of N steps

  Mean no. of Vertices $\langle V_n(N) \rangle \to$?

  Only $n = 1$ case (Open) walk, the result is known:

$$\boxed{\langle V_1(N) \rangle \simeq 2 \left[1 + \frac{1}{2} + \frac{1}{3} + \ldots + \frac{1}{N} \right] \sim 2 \log(N)} \quad \text{for large } N$$

(Baxter, '61)

- Distributions of the perimeter, area of the convex hull?

- Non-Brownian paths $\to$ anomalous diffusion, e.g., Lévy flights, external potential ?

Open Questions

- For n planar random walks each of N steps

  Mean no. of Vertices $\langle V_n(N) \rangle \to$?

  Only $n = 1$ case (Open) walk, the result is known:

$$\boxed{\langle V_1(N) \rangle \simeq 2 \left[1 + \frac{1}{2} + \frac{1}{3} + \ldots + \frac{1}{N} \right] \sim 2 \log(N)} \quad \text{for large } N$$

(Baxter, '61)

- Distributions of the perimeter, area of the convex hull?

- Non-Brownian paths $\to$ anomalous diffusion, e.g., Lévy flights, external potential ?

- Effect of Interactions between trajectories on convex hull?

Open Questions

- For n planar random walks each of N steps

  Mean no. of Vertices $\langle V_n(N) \rangle \to$?

  Only $n = 1$ case (Open) walk, the result is known:

$$\boxed{\langle V_1(N) \rangle \simeq 2 \left[1 + \frac{1}{2} + \frac{1}{3} + \ldots + \frac{1}{N} \right] \sim 2 \log(N)} \quad \text{for large } N$$

(Baxter, '61)

- Distributions of the perimeter, area of the convex hull?

- Non-Brownian paths $\to$ anomalous diffusion, e.g., Lévy flights, external potential ?

- Effect of Interactions between trajectories on convex hull?