

# “Explosive Percolation”

S. N. Dorogovtsev

*University of Aveiro  
and  
Ioffe Institute, St. Petersburg*

R. A. da Costa, SND, A. V. Goltsev, and J. F. F. Mendes,  
“Explosive percolation” transition is actually continuous,

arXiv:1009.2534

D. Achlioptas, R. M. D'Souza, and J. Spencer,  
Science 323, 1453 (2009).



simulation: 512,000 nodes

ordinary  
percolation:

“explosive  
percolation”:

ordinary: —  
explosive: —

$$\min(s_1 s_2, s_3 s_4)$$




???


How can it be?:  
discontinuity coexisting with critical power-law  
distributions and scaling above and below this transition  
- ???

# Representative model of “explosive percolation”

ordinary  
percolation:




“explosive  
percolation” ( $m = 2$ ):




# Relation between models


A



B



Simulations  $2 \times 10^9$  nodes:



$$S \propto \delta^\beta$$

$$\beta \approx 0.0476 \approx 1/18, \delta = t - t_c$$

# Estimate

Suppose  $\beta = 1/18$  and  $N = 10^{18}$ .

The smallest time interval is  $1/N$ .

Then a single step from the percolation threshold gives

$$S \sim (10^{-18})^{1/18} \sim 0.1$$

So simulations are virtually useless.

We must study the **infinite** system.

# Distributions

$n(s)$  and  $P(s)$  are for clusters

$$P(s) = sn(s)/\langle s \rangle, \quad \sum_s P(s) = 1 - S$$

$Q(s)$  is for merging clusters,  $\sum_s Q(s) = 1 - S^2$

$$Q_{\text{cum}}(s) + S^2 = [P_{\text{cum}}(s) + S]^2$$

$$\begin{aligned} Q(s) &= [P_{\text{cum}}(s) + P_{\text{cum}}(s+1) + 2S]P(s) \\ &= [2 - 2P(1) - 2P(2) - \dots - 2P(s-1) - P(s)]P(s) \end{aligned}$$


# Equations

$$\frac{\partial P(s, t)}{\partial t} = s \sum_{u+v=s} Q(u, t)Q(v, t) - 2sQ(s, t)$$

exactly describe the evolution of the distributions in the full range of  $t$  for the infinite system.


We solved numerically  $10^6$  equations, which gives precise description of the distributions for  $s \leq 10^6$ .

$$s \leq 10^6$$



Fitting by the law  $S_0 + C\delta^\beta$  gives  $S_0 < 0.005$ .


$$s \leq 10^6$$



# Scaling functions


$$P(s, t) = s^{1-\tau} f(s\delta^{1/\sigma})$$

$$Q(s, t) = s^{3-2\tau} g(s\delta^{1/\sigma})$$



# Ordinary percolation: scaling functions

$$P(s) = s^{1-\tau} f(s\delta^{1/\sigma})$$



**Table:** Percolation thresholds, critical exponents, and fractal and upper critical dimensions ( $m = 2$ )

	$t_c$	$\beta$	$\tau$	$\sigma$	$\gamma_P$	$\gamma_Q$	$d_f$	$d_u$
Ordinary	1/2	1	5/2	1/2	1	—	4	6
Explosive	0.923207508(2)	0.0555(1)	2.04762(2)	0.857(3)	1.111(1)	1.0556(5)	2.333(1)	2.445(1)

$$\begin{aligned}\tau &= 1 + \beta/(1 + 3\beta), & \sigma &= 1/(1 + 3\beta), & \gamma_P &= 1 + 2\beta, \\ \gamma_Q &= 1 + \beta, & d_f &= 2(1 + 3\beta), & d_u &= 2(1 + 4\beta)\end{aligned}$$

$$t_c(\infty) - t_c(N) \sim N^{-2/d_u}$$

If the distributions are power-law at the critical point, then  $S \sim \delta^\beta$ .

Indeed, above  $t_c$ , we have

$$Q(s) \cong 2SP(s)$$

at large  $s$ .

So the equation for the asymptotics contains only  $P(s)$  and  $S(t)$ . This equation is very similar to that for ordinary percolation.

Using  $P(s, t_c) = f(0)s^{1-\tau}$  as an initial condition, we solve this equation and find critical exponents and scaling functions above  $t_c$ .

## Relation between $\tau$ and $t_c$

$$P(s = 1, t_c) \sum_s s^{1-\tau} \approx 1$$

$$P(s = 1, t) = \frac{2}{1 + e^{4t}}$$

$$\frac{2}{1 + e^{4t_c}} \zeta(\tau - 1) \approx 1$$

Substituting  $t_c$  gives  $\tau - 2 \approx 0.05$ .

With increasing  $m$ ,  $t_c$  approaches 1 and  $\beta$  rapidly decreases with  $m$ , but the transition remains continuous.

# Conclusion

There is no explosion in “explosive percolation”.

# Lectures on Complex Networks

