Planar maps and continued fractions

Jérémie Bouttier, Emmanuel Guitter

Institut de Physique Théorique, CEA Saclay

Nordita – 3 November 2010

・ロト ・回ト ・ヨト

- ∢ ⊒ →

Maps: graphs embedded in surfaces (sphere in planar case) considered up to deformation (\Rightarrow finite number of maps with *E* edges) a.k.a. planar diagrams, fatgraphs, dynamical random tessellations...



イロン イヨン イヨン イヨン

Maps: graphs embedded in surfaces (sphere in planar case) considered up to deformation (\Rightarrow finite number of maps with *E* edges) a.k.a. planar diagrams, fatgraphs, dynamical random tessellations...



Motivations

- combinatorics [Tutte 1963]
- large N expansion of matrix integrals [Brézin-Itzykson-Parisi-Zuber 1979]
- 2D quantum gravity
- statistical physics on dynamical random surfaces
- probability theory: "Brownian map", connection with conformally-invariant processes

イロト イヨト イヨト イヨト

General model:

Each face of valency k comes with fugacity g_k :

$$Z = \sum_{\text{maps}} \prod_{k \ge 1} g_k^{\#\{k-\text{valent faces}\}}$$

Simple models: triangulations (resp. quadrangulations)

$$g_{k} = \begin{cases} g \text{ for } k = 3 \quad (\text{resp. } k = 4) \\ 0 \text{ otherwise} \end{cases} \qquad \qquad Z = \sum_{\substack{(\text{tri}|\text{quadr})\text{-}\\ \text{angulations}}} g^{\text{"area"}}$$

イロン イヨン イヨン イヨン

General model:

Each face of valency k comes with fugacity g_k :

$$Z = \sum_{\mathsf{maps}} \prod_{k \geq 1} g_k^{\#\{k-\mathsf{valent faces}\}}$$

Simple models: triangulations (resp. quadrangulations)

$$g_{k} = \begin{cases} g \text{ for } k = 3 \quad (\text{resp. } k = 4) \\ 0 \text{ otherwise} \end{cases} \qquad \qquad Z = \sum_{\substack{(\text{tri}|\text{quadr})-\\ \text{angulations}}} g^{\text{"area"}}$$

Here no extra "matter" degrees of freedom.

・ロン ・回と ・ヨン ・ヨン

Outline

1 First problem: maps with a boundary (review)

2 Second problem: maps with two points at given distance

・ロン ・回と ・ヨン・

Computing the partition function is an enumeration problem. It is simpler to count rooted maps



・ロト ・回ト ・ヨト

-≣->

Computing the partition function is an enumeration problem. It is simpler to count rooted maps with fixed root degree n



・ロト ・回ト ・ヨト

Computing the partition function is an enumeration problem. It is simpler to count rooted maps with fixed root degree n



・ロト ・回ト ・ヨト

< ≣⇒

Computing the partition function is an enumeration problem. It is simpler to count rooted maps with fixed root degree n i.e compute their generating function

 $F_n \equiv F_n(\{g_k\}_{k\geq 1}) = \frac{\partial Z}{\partial g_n}$ (w/o weight g_n for the root face).



・ロト ・回ト ・ヨト

Computing the partition function is an enumeration problem. It is simpler to count rooted maps with fixed root degree n i.e compute their generating function

 $F_n \equiv F_n(\{g_k\}_{k \ge 1}) = \frac{\partial Z}{\partial g_n}$ (w/o weight g_n for the root face).

 $F(z) = 1 + \sum_{n=1}^{\infty} F_n z^n$ is the disk amplitude.



イロン イヨン イヨン イヨン

Connection with matrix models

Consider a random $N \times N$ Hermitian matrix M with measure

$$dM \exp N\left(-\frac{Tr\,M^2}{2} + \sum_{k\geq 1} g_k \frac{Tr\,M^k}{k}\right)$$

then we have informally

$$F_n = \lim_{N \to \infty} \frac{1}{N} \langle Tr M^n \rangle$$

$$F(z) = \lim_{N \to \infty} \frac{1}{N} \langle Tr (1 - zM)^{-1} \rangle \qquad \text{(resolvent)}$$

・ロン ・回 と ・ ヨ と ・ ヨ と

2

Tutte's equation (1968) a.k.a. loop equation

The F_n are fully determined by the quadratic equation

$$F_{n} = \sum_{i=0}^{n-2} F_{i}F_{n-2-i} + \sum_{k\geq 1} g_{k}F_{n+k-2}$$



< **₩** ► < **⇒** ►

Tutte's equation (1968) a.k.a. loop equation

The F_n are fully determined by the quadratic equation

$$F(z) = 1 + z^2 F(z)^2 + \sum_{k \ge 1} g_k z^{2-k} \left(F(z) - \sum_{j=0}^{k-2} z^j F_j \right)$$



< **₩** ► < **⇒** ►

Tutte's equation (1968) a.k.a. loop equation

The F_n are fully determined by the quadratic equation

$$F(z) = 1 + z^{2}F(z)^{2} + \sum_{k \ge 1} g_{k}z^{2-k}F(z) + P(z^{-1})$$



・ 同・ ・ ヨ・

Review of the solution of Tutte's equation

By the previous equation

$$F(z) = rac{1}{2z^2} \left(1 - \sum_{k \geq 1} g_k z^{2-k} \pm \sqrt{\Delta(z)}
ight)$$

イロト イヨト イヨト イヨト

Review of the solution of Tutte's equation

By the previous equation

$$F(z) = rac{1}{2z^2} \left(1 - \sum_{k \geq 1} g_k z^{2-k} \pm \sqrt{\Delta(z)}
ight)$$

By Brown's lemma/one-cut hypothesis

$$F(z) = \frac{1}{2z^2} \left(1 - \sum_{k \ge 1} g_k z^{2-k} - \Gamma(z^{-1}) \sqrt{1 + \kappa_1 z + \kappa_2 z^2} \right)$$

with $\Gamma(z^{-1})$ a polynomial or power series in z^{-1} .

(4回) (4回) (日)

Review of the solution of Tutte's equation

By the previous equation

$$F(z) = rac{1}{2z^2} \left(1 - \sum_{k \geq 1} g_k z^{2-k} \pm \sqrt{\Delta(z)}
ight)$$

By Brown's lemma/one-cut hypothesis

$$F(z) = \frac{1}{2z^2} \left(1 - \sum_{k \ge 1} g_k z^{2-k} - \Gamma(z^{-1}) \sqrt{1 + \kappa_1 z + \kappa_2 z^2} \right)$$

with $\Gamma(z^{-1})$ a polynomial or power series in z^{-1} .

But F(z) contains only nonnegative powers of z! This constraint allows to deduce explicit expressions for $\Gamma(z^{-1})$, κ_1 , κ_2 .

Example: quadrangulations

For
$$g_k = \begin{cases} g \text{ for } k = 4\\ 0 \text{ otherwise} \end{cases}$$
 this method leads to
$$F_{2n} = \sum_{a=0}^{\infty} \frac{(2n)!}{n!(n-1)!} \frac{(2a+n-1)!}{a!(a+n+1)!} (3g)^a \qquad F_{2n+1} = 0$$

・ロト ・回 ト ・ヨト ・ヨト

æ

For
$$g_k = \begin{cases} g \text{ for } k = 4\\ 0 \text{ otherwise} \end{cases}$$
 this method leads to
$$F_{2n} = \sum_{a=0}^{\infty} \frac{(2n)!}{n!(n-1)!} \frac{(2a+n-1)!}{a!(a+n+1)!} (3g)^a \qquad F_{2n+1} = 0$$

The sum converges for $g \le g_c = 1/12$. Similar expressions exist for triangulations (where $g_c = \sqrt{4/27}$). It is now easily to analyze the (well-known) critical behaviour:

- for fixed finite n, as $g \to g_c$, $\partial F_n/\partial g$ and $\partial^2 Z/\partial g^2$ have a square-root singularity (" $\gamma_{\text{string}} = -1/2$ "),
- the relevant scaling is $n \propto 1/\sqrt{g_c g}$: the dominant singular term of F_n corresponds to the universal disk amplitude of pure gravity.

General combinatorial structure of the solution

$$F(z) = \frac{1}{2z^2} \left(1 - \sum_{k \ge 1} g_k z^{2-k} - \Gamma(z^{-1}) \sqrt{1 + \kappa_1 z + \kappa_2 z^2} \right) \quad (1)$$

・ロト ・回ト ・ヨト ・ヨト

æ

$$F(z) = \frac{1}{2z^2} \left(1 - \sum_{k \ge 1} g_k z^{2-k} - \Gamma(z^{-1}) \sqrt{1 + \kappa_1 z + \kappa_2 z^2} \right) \quad (1)$$

Trick: replace the unknowns κ_1, κ_2 by R, S with

$$\kappa(z) \equiv 1 + \kappa_1 z + \kappa_2 z^2 = (1 - Sz)^2 - 4Rz^2$$

・ロト ・回ト ・ヨト ・ヨト

$$F(z) = \frac{1}{2z^2} \left(1 - \sum_{k \ge 1} g_k z^{2-k} - \Gamma(z^{-1}) \sqrt{1 + \kappa_1 z + \kappa_2 z^2} \right) \quad (1)$$

Trick: replace the unknowns κ_1, κ_2 by R, S with

$$\kappa(z) \equiv 1 + \kappa_1 z + \kappa_2 z^2 = (1 - Sz)^2 - 4Rz^2$$

then

$$\sqrt{\kappa(z)} = 1 - Sz - 2Rz^2 \sum_{n=0}^{\infty} P^+(n; R, S) z^n$$

・ロト ・回ト ・ヨト ・ヨト

$$F(z) = \frac{1}{2z^2} \left(1 - \sum_{k \ge 1} g_k z^{2-k} - \Gamma(z^{-1}) \sqrt{1 + \kappa_1 z + \kappa_2 z^2} \right) \quad (1)$$

Trick: replace the unknowns κ_1, κ_2 by R, S with

$$\kappa(z) \equiv 1 + \kappa_1 z + \kappa_2 z^2 = (1 - Sz)^2 - 4Rz^2$$

then

$$\sqrt{\kappa(z)} = 1 - Sz - 2Rz^2 \sum_{n=0}^{\infty} P^+(n; R, S) z^n$$

 $P^+(n; R, S)$ is the generating function for Motzkin paths of length *n*, with weight *R* (resp. *S*) per down-step (resp. level-step).

$$\begin{array}{c} \checkmark R \\ \checkmark S \\ (0,0) \\ P^{+}(n;R,S) \end{array}$$

(1) immediately yields

$$F_n = R \sum_{q \ge 0} \gamma_q P^+(n+q;R,S)$$
⁽²⁾

The only dependence in n is via the path length!

・ 回 と ・ ヨ と ・ ヨ と

(1) immediately yields

$$F_n = R \sum_{q \ge 0} \gamma_q P^+(n+q;R,S)$$
⁽²⁾

The only dependence in n is via the path length!

By now writing that (1) (divided by $\sqrt{\kappa(z)}$) contains no negative powers in z and that its constant term is 1, we may obtain:

- algebraic equations determining the "master unknowns" R, S
- expressions for the γ_q in terms of R, S.

・ 同 ト ・ ヨ ト ・ ヨ ト

(1) immediately yields

$$F_n = R \sum_{q \ge 0} \gamma_q P^+(n+q;R,S)$$
⁽²⁾

The only dependence in *n* is via the path length! By now writing that (1) (divided by $\sqrt{\kappa(z)}$) contains no negative powers in *z* and that its constant term is 1, we may obtain:

- algebraic equations determining the "master unknowns" R, S
- expressions for the γ_q in terms of R, S.

Remark: these may also be given a combinatorial interpretation via

$$1/\sqrt{\kappa(z)} = \sum_{n=0}^{\infty} P(n; R, S) z^n$$



Summary/conclusion on the first problem

- Maps with a boundary can be enumerated effectively via Tutte's equation.
- A remarkable combinatorial/algebraic structure related to the physical one-cut hypothesis.
- *F*(*z*) is a master function in terms of which generating functions for maps with several boundaries and of higher genus ("global observables") can be expressed.
- Generalizations to models with matter are known.

・ロン ・回と ・ヨン ・ヨン

Outline



2 Second problem: maps with two points at given distance

イロン イヨン イヨン イヨン



Geodesic/graph distance: minimal number of edges connecting two given vertices (i.e each edge has length 1)

What are the metric properties of large random maps?

・ロト ・回ト ・ヨト



Simple observable: the distance-dependent two-point function [Ambjørn-Watabiki 1996] is the generating function for maps with two marked points at given distance. Computing it is again an enumeration problem!

Probabilistic interpretation: it encodes the distribution of distances between two uniformly chosen random random points.

A (1) < A (1) </p>

By a transfer matrix approach, Ambjørn and Watabiki successfully predicted the universal scaling form of the two-point function for pure gravity.

A (1) > (1) > (1)

By a transfer matrix approach, Ambjørn and Watabiki successfully predicted the universal scaling form of the two-point function for pure gravity.

Scaling: distance
$$\propto (g_c-g)^{-1/4} \propto ($$
area $)^{1/4}$

In a canonical ensemble (maps of fixed area), the rescaled distance between two uniform random points admits a limiting distribution:



$$\sim d^3 ext{ for } d o 0$$

 $\sim e^{-Cd^{4/3}} ext{ for } d o \infty$

 \sim

An exact discrete expression whose scaling form agrees with the Ambjørn-Watabiki prediction was found for quadrangulations and, more generally, maps with even face valencies. [B., Di Francesco, Guitter 2003]

< A > < 3

An exact discrete expression whose scaling form agrees with the Ambjørn-Watabiki prediction was found for quadrangulations and, more generally, maps with even face valencies. [B., Di Francesco, Guitter 2003]

Ingredients:

- coding of maps by trees (Schaeffer's bijection and generalizations)
- identification of the two-point function with tree g.f.
- equation following from recursive decomposition of such trees
- guess of the solution!

・ロト ・ 日 ・ ・ ヨ ト

The discrete two-point function is the solution of the equation

$$R_n = 1 + gR_n(R_{n-1} + R_n + R_{n+1}) \qquad (n \ge 1, R_0 = 0)$$

The discrete two-point function is the solution of the equation

$$R_n = 1 + gR_n(R_{n-1} + R_n + R_{n+1})$$
 $(n \ge 1, R_0 = 0)$

Explicit solution

$$R_n = R \frac{u_n u_{n+3}}{u_{n+1} u_{n+2}} \tag{3}$$

・ロト ・回ト ・ヨト

∃ >

$$R = 1 + 3gR^2$$
 $u_n = 1 - x^n$ $x + \frac{1}{x} + 1 = \frac{1}{gR^2}$

The discrete two-point function is the solution of the equation

$$R_n = 1 + gR_n(R_{n-1} + R_n + R_{n+1})$$
 $(n \ge 1, R_0 = 0)$

Explicit solution $R_{n} = R \frac{u_{n}u_{n+3}}{u_{n+1}u_{n+2}}$ (3) $R = 1 + 3gR^{2} \qquad u_{n} = 1 - x^{n} \qquad x + \frac{1}{x} + 1 = \frac{1}{gR^{2}}$

There are also equations with explicit solutions in more general cases! The form (3) still holds (but u_n gets more complicated). Why?

◆□ > ◆□ > ◆臣 > ◆臣 > ○

New approach [B., Guitter 2010]

The two-point function is encoded in the continued fraction expansion of the disk amplitude F(z)!

• Maps with even face valencies: Stieljes fraction

$$F(z) \equiv \sum_{n=0}^{\infty} F_{2n} z^{2n} = \frac{1}{1 - \frac{R_1 z^2}{1 - \frac{R_2 z^2}{1 - \cdots}}}$$

• Maps with arbitrary face valencies: Jacobi fraction

$$F(z) \equiv \sum_{n=0}^{\infty} F_n z^n = \frac{1}{1 - S_0 z - \frac{R_1 z^2}{1 - S_1 z - \frac{R_2 z^2}{1 - \cdots}}}$$
(4)

Elements of the proof:

• the combinatorial theory of continued fractions [Flajolet 1980]

Combinatorial equivalent of (4)

 F_n is equal to the generating function for Motzkin paths of length n, with weight R_m (resp. S_m) per down-step (resp. level-step) starting at height m.

イロト イヨト イヨト イヨト

Elements of the proof:

• the combinatorial theory of continued fractions [Flajolet 1980]

Combinatorial equivalent of (4)

 F_n is equal to the generating function for Motzkin paths of length n, with weight R_m (resp. S_m) per down-step (resp. level-step) starting at height m.

• a suitable decomposition of maps with a boundary (via trees or "slices"): Motzkin paths code the distances from the origin to the vertices incident to the root face.



・ロン ・回 と ・ヨン ・ヨン

æ

Via Hankel determinants:

$$R_n = \frac{H_n H_{n-2}}{H_{n-1}^2} \qquad H_n = \det_{0 \le i, j \le n} F_{i+j}$$
$$S_n = \frac{\tilde{H}_n}{H_n} - \frac{\tilde{H}_{n-1}}{H_{n-1}} \qquad \tilde{H}_n = \det_{0 \le i, j \le n} F_{i+j+\delta_{j,n}}$$

・ロン ・回と ・ヨン ・ヨン

æ

Via Hankel determinants:

$$R_n = \frac{H_n H_{n-2}}{H_{n-1}^2} \qquad H_n = \det_{0 \le i,j \le n} F_{i+j}$$
$$S_n = \frac{\tilde{H}_n}{H_n} - \frac{\tilde{H}_{n-1}}{H_{n-1}} \qquad \tilde{H}_n = \det_{0 \le i,j \le n} F_{i+j+\delta_{j,n}}$$

Even face valencies: $S_n = \tilde{H}_n = 0$, H_n factorizes as:

$$H_{2n} = h_n^{(0)} h_{n-1}^{(1)}$$
 $H_{2n+1} = h_n^{(0)} h_n^{(1)}$ $h_n^{(e)} = \det_{0 \le i,j \le n} F_{2i+2j+2e}$

This essentially explains the form (3).

・ロン ・回と ・ヨン・

Via Hankel determinants:

$$R_n = \frac{H_n H_{n-2}}{H_{n-1}^2} \qquad H_n = \det_{0 \le i,j \le n} F_{i+j}$$
$$S_n = \frac{\tilde{H}_n}{H_n} - \frac{\tilde{H}_{n-1}}{H_{n-1}} \qquad \tilde{H}_n = \det_{0 \le i,j \le n} F_{i+j+\delta_{j,n}}$$

Even face valencies: $S_n = \tilde{H}_n = 0$, H_n factorizes as:

$$H_{2n} = h_n^{(0)} h_{n-1}^{(1)}$$
 $H_{2n+1} = h_n^{(0)} h_n^{(1)}$ $h_n^{(e)} = \det_{0 \le i,j \le n} F_{2i+2j+2e}$

This essentially explains the form (3).

These relations hold in the general theory of continued fractions. In our map model, the specific form of F_n lead to specific Hankel determinants, which are symplectic Schur functions $\operatorname{sp}_{2p}(\lambda, \mathbf{x})$.

The general formula for F_n is

$$F_n = \sum_{q=0}^{p} A_q P^+(n+q)$$

Substituting into the Hankel determinant

1

 $P_{k}(q;R,S) - P(q;R,S)$

q

 $P^{\dagger}(i+j+q;R,S)$

-2-1012

◆□ > ◆□ > ◆臣 > ◆臣 > ○

æ

 $P_{l}^{\dagger}(j;R,S)$

 $P_k^+(i;R,S)$

Remark

We make use of *two* different formulas for F_n involving Motzkin paths:

- as a sum (2) over Motzkin paths of variable length
 n,..., n + p and height-independant weights R, S per step
- as a sum (4) over Motzkin paths of fixed length *n* and height-dependant weights R_m, S_m per step

イロン イヨン イヨン イヨン

Remark

We make use of *two* different formulas for F_n involving Motzkin paths:

- as a sum (2) over Motzkin paths of variable length
 n,..., n + p and height-independant weights R, S per step
- as a sum (4) over Motzkin paths of fixed length *n* and height-dependant weights R_m, S_m per step

Caveat

The expression involving Schur functions assumes that face valencies are bounded: $g_k = 0$ for k > p + 2. H_n may then be rewritten as a $p \times p$ determinant (rather than $(n+1) \times (n+1)$), easier to study in the limit of large distance n.

・ロン ・回 と ・ ヨ と ・ ヨ と

Example & combinatorial interpretation: triangulations

Suppose that $g_k = 0$ for $k \neq 3$ (faces are triangles), i.e p = 1:

$$F_n = A_0 P^+(n; R, S) + A_1 P^+(n+1; R, S)$$

・ロト ・回ト ・ヨト ・ヨト

Example & combinatorial interpretation: triangulations

Suppose that $g_k = 0$ for $k \neq 3$ (faces are triangles), i.e p = 1:

$$F_n = A_0 P^+(n; R, S) + A_1 P^+(n+1; R, S)$$



(ロ) (同) (E) (E) (E)

Example & combinatorial interpretation: triangulations

Suppose that $g_k = 0$ for $k \neq 3$ (faces are triangles), i.e p = 1:

$$F_n = A_0 P^+(n; R, S) + A_1 P^+(n+1; R, S)$$



 F_{i+j} can be interpreted as paths on a weighted graph. By the Lindström-Gessel-Viennot lemma, the determinant H_n counts configurations of non-intersecting lattice paths on this graph.

Such configurations of non-intersecting lattice paths are highly constrained and, actually, in bijection with configurations of 1D dimers.



・ロト ・回ト ・ヨト ・ヨト

Counting 1D dimer configurations is easy, we obtain

$$H_n \propto rac{1}{(1+y)^{n+1}} rac{1-y^{n+2}}{1-y}$$

with y related to the dimer weight $-g_3^2 R^3$ by

$$y + \frac{1}{y} + 2 = \frac{1}{g_3^2 R^3}.$$

It yields the simple formula

$$R_n = R \frac{(1-y^n)(1-y^{n+2})}{(1-y^{n+1})^2}$$

and similarly

$$S_n = S - g_3 R^2 y^n \frac{(1-y)(1-y^2)}{(1-y^{n+1})(1-y^{n+2})}$$

・ロン ・回と ・ヨン ・ヨン

Conclusion and outlook

- We have shown that the disk amplitude and the two-point function are encoded in the same function F(z).
- Our results are purely discrete. One may now turn to asymptotic analysis. The generic behaviour is pure gravity ("Brownian map").
- Possible directions:
 - Connections with orthogonal polynomials and matrix models
 - Other distance-related observables (not so many known! radius, three-point function, numbers of geodesics...)
 - Generalizations to models with matter
 - Maps with large faces?

References:

- J. Bouttier, P. Di Francesco and E. Guitter, Nucl.Phys. B663 (2003) 535-567, arXiv:cond-mat/0303272,
- J. Bouttier and E. Guitter, arXiv:1007.0419.

Summary: the two facets of F(z)



・ロン ・回 と ・ ヨン ・ ヨン

3

Summary: the two facets of F(z)



・ロン ・回 と ・ ヨン ・ ヨン