

Discussions

- now
- tomorrow morning: galactic issues
- tomorrow afternoon: extragalactic issues
- Friday morning: astroparticle physics, fundamental physics, and all the rest





Discussion I

• What might be "first-day" studies, high-priority targets which need to be attacked immediately?



- What might be "first-day" studies, high-priority targets which need to be attacked immediately?
- CTA might, in a pessimistic scenario, not have overlap with LAT, while HESS-II still has. What strategic steps can be taken to exploit this optimally?



- What might be "first-day" studies, high-priority targets which need to be attacked immediately?
- CTA might, in a pessimistic scenario, not have overlap with LAT, while HESS-II still has. What strategic steps can be taken to exploit this optimally?
- The Crab Nebula was believed to be the standard candle of gamma-ray astronomy for many years. What are the implications of the flares observed in the HE regime?



- What might be "first-day" studies, high-priority targets which need to be attacked immediately?
- CTA might, in a pessimistic scenario, not have overlap with LAT, while HESS-II still has. What strategic steps can be taken to exploit this optimally?
- The Crab Nebula was believed to be the standard candle of gamma-ray astronomy for many years. What are the implications of the flares observed in the HE regime?
 - what can HESS-II add in studying and understanding the flares?



- What might be "first-day" studies, high-priority targets which need to be attacked immediately?
- CTA might, in a pessimistic scenario, not have overlap with LAT, while HESS-II still has. What strategic steps can be taken to exploit this optimally?
- The Crab Nebula was believed to be the standard candle of gamma-ray astronomy for many years. What are the implications of the flares observed in the HE regime?
 - what can HESS-II add in studying and understanding the flares?
 - are there implications from the (current understanding of the) flares that imply future studies with HESS-II that were not possible with HESS before?



- What might be "first-day" studies, high-priority targets which need to be attacked immediately?
- CTA might, in a pessimistic scenario, not have overlap with LAT, while HESS-II still has. What strategic steps can be taken to exploit this optimally?
- The Crab Nebula was believed to be the standard candle of gamma-ray astronomy for many years. What are the implications of the flares observed in the HE regime?
 - what can HESS-II add in studying and understanding the flares?
 - are there implications from the (current understanding of the) flares that imply future studies with HESS-II that were not possible with HESS before?
- The current limitations of DM searches with HE/VHE gamma-ray experiments are coming from the modeling of the DM content of the sources. What are the prospects for the coming years?



- What might be "first-day" studies, high-priority targets which need to be attacked immediately?
- CTA might, in a pessimistic scenario, not have overlap with LAT, while HESS-II still has. What strategic steps can be taken to exploit this optimally?
- The Crab Nebula was believed to be the standard candle of gamma-ray astronomy for many years. What are the implications of the flares observed in the HE regime?
 - what can HESS-II add in studying and understanding the flares?
 - are there implications from the (current understanding of the) flares that imply future studies with HESS-II that were not possible with HESS before?
- The current limitations of DM searches with HE/VHE gamma-ray experiments are coming from the modeling of the DM content of the sources. What are the prospects for the coming years?
- In the view of the recent results from collider experiments, and in particular the ² discovery of a Higgs-like particle at 125 GeV, how natural is SUSY nowadays? What are the implications of this discovery for extra-dimensions models?





Discussion II

• Now that we have more and more evidence of hadrons acceleration in SNR, how far are we to conclude on the primary origin of galactic CRs?



- Now that we have more and more evidence of hadrons acceleration in SNR, how far are we to conclude on the primary origin of galactic CRs?
- In case of binary system PSR B1259-63, a huge flaring episode was detected by Fermi-LAT "after" the periastron passage, but not by IACT experiments, whereas the baseline emission was observed in the two energy regimes. What does this tell us on the origin of emission of the HE component in these objects? What can we hope for with HESS2 observations?





Discussion III

• How well do we know the EBL and what are the prospects to improve this knowledge in the coming years? What are the implications of these uncertainties for observations of distant gamma-ray sources?



- How well do we know the EBL and what are the prospects to improve this knowledge in the coming years? What are the implications of these uncertainties for observations of distant gamma-ray sources?
- What would be the implications of observing one GRB with an IACT? Since we already know that different population of GRBs exist, how many GRBs measurements in the VHE domain would be necessary to understand the underlying physics and put some guesses on their origin?



- How well do we know the EBL and what are the prospects to improve this knowledge in the coming years? What are the implications of these uncertainties for observations of distant gamma-ray sources?
- What would be the implications of observing one GRB with an IACT? Since we already know that different population of GRBs exist, how many GRBs measurements in the VHE domain would be necessary to understand the underlying physics and put some guesses on their origin?
- What can we learn from being able to access lower energies concerning the sources intrinsically?



- How well do we know the EBL and what are the prospects to improve this knowledge in the coming years? What are the implications of these uncertainties for observations of distant gamma-ray sources?
- What would be the implications of observing one GRB with an IACT? Since we already know that different population of GRBs exist, how many GRBs measurements in the VHE domain would be necessary to understand the underlying physics and put some guesses on their origin?
- What can we learn from being able to access lower energies concerning the sources intrinsically?