



© P-M Heden

**Extra-galactic background Light – Dieter Horns (Univ. Hamburg)**



Daniel López  
Observatorio del Teide, IAC


Workshop Stockholm

Dieter Horns  
<http://www-hess.desy.de>

May 13, 2014



# The sky is bright - the universe is not..




Leinert et al. 1998




# The Extra-galactic background light in a box

- Counting photons in a  $1 \text{ m}^3$  box:

$4 \times 10^8$  Cosmic-microwave background photons



Somewhere in Extra-galactic space






# The Extra-galactic background light in a box

- Counting photons in a  $1 \text{ m}^3$  box:

4  $\times 10^8$  Cosmic microwave background photons  
 $\sim 10^6$  Cosmic infra-red background photons



Somewhere in Extra-galactic space




View from Earth




# The Extra-galactic background light in a box

- Counting photons in a  $1 \text{ m}^3$  box:




4  $\times 10^8$  Cosmic microwave background photons  
 $\sim 10^6$  Cosmic infra-red background photons  
 $\sim 10^4$  Cosmic optical photons




Somewhere in Extra-galactic space



# Dissecting the EBL with the Hubble telescope



Resolved Sources  
(galaxies):




[Madau&Pozzetti: Astro-ph/9907315]



# Dissecting the EBL with the Hubble telescope



Unresolved Sources  
(too faint)






# Dissecting the EBL with the Hubble telescope



Unresolved and  
unknown Sources  
(too faint)





# Dissecting the EBL with the Hubble telescope




Additional contribution  
(particle decays,...)

The Unknown  
Unknowns....

Cambresy et al. 2001  
[astro-ph/0103078]



# Dissecting the EBL with the Hubble telescope




Background fluctuations  
Unresolved contribution  
 $<1 \text{ nW}/(\text{m}^2\text{srad})$




# NIR / X-ray excess clustering at degree-scale


Yue et al. 2013 [arXiv:1305.5177]



- No correlation with HST maps (Kashlinsky et al. 2007) → high z
- Field stars? (Cooray et al. 2013)
- Correlation with X-ray counts (Capelluti et al. 2013, Helgason 2012, 2014) → Direct black hole collapse (Yue et al. 2013)
- 0.5-2 nW/m<sup>2</sup>srad: Limiting DM star contribution (Maurer et al. 2012)



# Models (one slide only)



Different models &  
different methods →  
Converging results!

The simplest type of model (Dwek et al. 1998,  
Kneiske et al. 2002)

$$P_\nu(z) = \nu I_\nu(z) = \nu \frac{c}{4\pi} \int_z^{z_m} \mathcal{E}_\nu(z') \left| \frac{dt'}{dz'} \right| dz'$$


EBL

$$\mathcal{E}_\nu(z) = \int_z^{z_m} L_\nu(t(z) - t(z')) \dot{\rho}_*(z') \left| \frac{dt'}{dz'} \right| dz'$$

Emissivity

Stellar population spectra (SPS)


Star formation rate density (SFRD)






# Dark stars' contribution to the EBL

Synthetic model spectra for DS




Plug in your favorite model  
and estimate the contribution



$$\begin{aligned}
 (vI_v)_{\text{max}} = & 2 \times 10^{-3} \text{ nW m}^{-2} \text{ sr}^{-1} \times \left( \frac{\text{---DS}}{10^7 \text{ years}} \right) \\
 & \times \left( \frac{\text{SFR}_{\text{Norm}}}{10^{-5}} \right) \times \left( \frac{\text{LMR}}{10^3 L_{\odot} / M_{\odot}} \right) \times \left( \frac{z_{\text{min}}}{10} \right)^{-2.5}
 \end{aligned}$$



## Absolute Measurements of the EBL



Dwek & Krennrich  
[arXiv:1209.4661]



# Transparency of the Universe to gamma-rays

Astrophysics

Extra-galactic  
Background light  
(EBL)

Astroparticle  
Physics

Propagation of  
Extra-galactic  
Gamma-rays

Particle Physics


Peccei-Quinn symmetry  
And axion-like particles

→ A. Ringwald



# Very high energy gamma-ray observations

High Energy  
Stereoscopic System  
(H.E.S.S.)




Excellent review: Hillas (2013)




# Gamma-ray attenuation via pair-production

[Nikishov 1962, Jelley 1966, Gould&Schréder 1966]



$$\tau_\gamma(E, z_0) = \int_0^{z_0} d\ell(z) \int_{-1}^{+1} d\mu \frac{1-\mu}{2} \int_{\epsilon_{\text{thr}}}^{\infty} d\epsilon' n_{\text{EBL}}(\epsilon', z) \sigma_{\gamma\gamma}(E, \epsilon', \mu)$$


$$\frac{dN_{\text{obs}}}{dE} = \frac{dN_{\text{int}}}{dE} \times \exp[-\tau_\gamma(E, z_0)]$$






# Gamma-ray attenuation via pair-production

[Nikishov 1962, Jelley 1966, Gould&Schréder 1966]



$$\tau_\gamma(E, z_0) = \int_0^{z_0} d\ell(z) \int_{-1}^{+1} d\mu \frac{1-\mu}{2} \int_{\epsilon_{\text{thr}}}^{\infty} d\epsilon' n_{\text{EBL}}(\epsilon', z) \sigma_{\gamma\gamma}(E, \epsilon', \mu)$$


$$\frac{dN_{\text{obs}}}{dE} = \frac{dN_{\text{int}}}{dE} \times \exp[-\tau_\gamma(E, z_0)]$$






# Gamma-ray attenuation via pair-production

[Nikishov 1962, Jelley 1966, Gould&Schréder 1966]



$$\tau_\gamma(E, z_0) = \int_0^{z_0} d\ell(z) \int_{-1}^{+1} d\mu \frac{1-\mu}{2} \int_{\epsilon_{\text{thr}}}^{\infty} d\epsilon' n_{\text{EBL}}(\epsilon', z) \sigma_{\gamma\gamma}(E, \epsilon', \mu)$$



$$\frac{dN_{\text{obs}}}{dE} = \frac{dN_{\text{int}}}{dE} \times \exp [-\tau_\gamma(E, z_0)]$$






# Measuring the EBL with VHE gamma-ray attenuation

Raue & Mazin 2010  
[arXiv:1005.1196]





# First detection of the EBL (1-5 $\mu\text{m}$ ) at VHE (0.2-20 TeV) energies!!



PKS 2005–489 at VHE: four years of monitoring with HESS and simultaneous multi-wavelength observations

Discovery of VHE  $\gamma$ -rays from the distant BL Lacertae 1ES 0347-121\*

New constraints on the mid-IR EBL from the HESS discovery of VHE  $\gamma$ -rays from 1ES 0229+200

**75 000  $\gamma$ -rays from the seven brightest blazars, with  $0.03 < z < 0.19$ , collected during 400 hours with H.E.S.S.**

Source	$z$	$N_{\gamma}$	$E_{\min} - E_{\max}$ [TeV]
Mrk 421 (1)	0.031	3381	0.95 – 41
Mrk 421 (2)	0.031	5548	0.95 – 37
Mrk 421 (3)	0.031	5156	0.95 – 45
PKS 2005-489 (1)	0.071	1540	0.16 – 37
PKS 2005-489 (2)	0.071	910	0.18 – 25
PKS 2155-304 (2008)	0.116	5279	0.13 – 19
PKS 2155-304 (1)	0.116	3499	0.13 – 5.7
PKS 2155-304 (2)	0.116	3470	0.13 – 9.3
PKS 2155-304 (3)	0.116	9555	0.13 – 14
PKS 2155-304 (4)	0.116	4606	0.18 – 4.6
PKS 2155-304 (5)	0.116	11901	0.13 – 5.7
PKS 2155-304 (6)	0.116	6494	0.15 – 5.7
PKS 2155-304 (7)	0.116	8253	0.20 – 7.6
1ES 0229+200	0.14	670	0.29 – 25
H 2356-309	0.165	1642	0.11 – 34
1ES 1101-232	0.186	1268	0.12 – 23
1ES 0347-121	0.188	604	0.13 – 11

Data sets on highly significant sources were divided and sorted by flux level

Biteau,  
H.E.S.S coll. (2013)  
[arXiv:1212.3459]






# First detection of the EBL (1-5 $\mu\text{m}$ ) at VHE (0.2-20 TeV) energies!!

H.E.S.S coll. (2013)  
[arXiv:1212.3459]

Fit of normalization:


$$\phi(E, z) = \Phi(E) \cdot \exp(-\alpha \tau(E, z))$$






# Detection of the EBL at high energies

Fermi-LAT coll. Science 2012 [arXiv:1211.1671]




Similar results  
Gong&Cooray [arXiv:1305.5249]





# Consistency of the EBL-measurements





# Hidden treasures?




- EBL measurement leaves little room for additional components (QSO at UV, first stars, decaying dark matter, ...)
- In fact, the measured EBL falls below the lower limit from Galaxy counts



# Indications for modified optical depth

Adapted from Dwek&Krennrich 2012


- De Angelis et al. 2009, 2011, 2013:  
The observed TeV spectra are too hard.





# Indications for modified optical depth


- De Angelis et al. 2009, 2011, 2013:  
The observed TeV spectra are too hard.
- DH and Meyer 2012 [arXiv:1201.4711]:  
The attenuation of VHE spectra at  $\tau > 2$  is  
too small (at  $\sim 4 \sigma$ )





# Indications for modified optical depth


- De Angelis et al. 2009, 2011, 2013:  
The observed TeV spectra are too hard.
- DH and Meyer 2012 [arXiv:1201.4711]:  
The attenuation of VHE spectra at  $\tau > 2$  is  
too small (at  $\sim 4 \sigma$ )





# Indications for modified optical depth


- De Angelis et al. 2009, 2011, 2013:  
The observed TeV spectra are too hard.
- DH and Meyer 2012 [arXiv:1201.4711]:  
The attenuation of VHE spectra at  $\tau > 2$  is  
too small (at  $\sim 4 \sigma$ )





# Indications for modified optical depth


- De Angelis et al. 2009, 2011, 2013:  
The observed TeV spectra are too hard.
- DH and Meyer 2012 [arXiv:1201.4711]:  
The attenuation of VHE spectra at  $\tau > 2$  is  
too small (at  $\sim 4 \sigma$ )
- Meyer 2013 (desy-thesis 2013-33),  
DH and Meyer 2013 [arXiv:1309.3846]:  
Fermi-LAT photon ( $\sim 3.5\sigma$ ) excess at  $\tau > 2$   
(similar: [arXiv:1207.1962])





# Indications for modified optical depth

- De Angelis et al. 2009, 2011, 2013:  
The observed TeV spectra are too hard.
- DH and Meyer 2012 [arXiv:1201.4711]:  
The attenuation of VHE spectra at  $\tau > 2$  is  
too small (at  $\sim 4 \sigma$ )
- Meyer 2013 (desy-thesis 2013-33),  
DH and Meyer 2013 [arXiv:1309.3846]:  
Fermi-LAT photon ( $\sim 3.5\sigma$ ) excess at  $\tau > 2$   
(similar: [arXiv:1207.1962])
- Very similar analysis of individual spectra  
(e.g. [arXiv:1308.380])





## Proposed ways to explain the transparency anomaly:

- Gamma-ray/cosmic-ray induced cascades → TeV emission is secondary nature (e.g., Essay & Kusenko 2010)
- Lorentz invariance violation (e.g., Jacob&Piran 2008, Shao&Ma 2010)
- Oscillations into hidden photons (e.g., Jaeckel&Ringwald 2010; Jaeckel 2013 for reviews)
- Oscillations into axion-like particles (e.g., Csaki et al. 2003, Jaeckel&Ringwald 2013, Carosi et al. 2013)



# Transparency of the Universe to gamma-rays

Astrophysics

Extra-galactic  
Background light  
(EBL)

Astroparticle  
Physics

Propagation of  
Extra-galactic  
Gamma-rays

Particle Physics

Peccei-Quinn symmetry  
And axion-like particles


→ A. Ringwald



## Peccei-Quinn symmetries $\leftrightarrow$ Axion & ALPs

- Additional symmetry to cure strong CP problem (Peccei&Quinn 77), leading „axion“ particle (Weinberg 78, Wilczek 78) with non-vanishing coupling to photons
- Generic  $U(1)_{\text{PQ}} \rightarrow \mathcal{L} \supset -\frac{g_{a\gamma}}{4} a F_{\mu\nu} \tilde{F}^{\mu\nu}$ 
  - Motivated by string compactifications („top bottom“)  $\rightarrow$  axiverse
  - Phenomenological solution to CP-Problem, Dark Matter, neutrino-masses, baryon asymmetry, stability of vacuum
- *Rich phenomenology through photon coupling of ALPs (Jaeckel&Ringwald 2010)*

(credit M.Meyer)




[e.g., De Angelis et al., 2007, 2011; Mirizzi et al., 2007; Simet et al., 2008;  
Sanchez-Condé et al., 2009; Horns et al. 2012; Tavecchio et al. 2012]



# Effect of gamma-ALPs mixing on propagation

1ES 0414+009


(DH, Maccione, Meyer, Mirizzi, Montanino, Roncadelli, arXiv: 1207.0776)





# Effect of gamma-ALPs mixing on propagation


(DH, Maccione, Meyer, Mirizzi, Montanino, Roncadelli, arXiv: 1207.0776)





# Effect of gamma-ALPs mixing on propagation


(DH, Maccione, Meyer, Mirizzi, Montanino, elli, arXiv: 1207.0776)







# Effect of gamma-ALPs mixing on propagation

(DH, Maccione, Meyer, Mirizzi, Montanino, Sillie, arXiv: 1207.0776)






# Lower limit on coupling $g_{a\gamma}$



- Coupling within reach of future helioscope IAXO (Irastorza et al. 2013), light-shining-through the wall ALPS-II (Bähre et al. [arXiv:1302.5647])
- Coupling consistent with anomalous cooling of WD (Isern et al. 2008)




# A closer look on PKS 1424+240





# A closer look on PKS 1424+240





# Summary

- Extra-galactic background light constrained by galaxy counts and direct measurements:  $10..30 \text{ nW}/(\text{m}^2 \text{ srad})$  in the optical/NIR
- Detection of absorption feature with Fermi-LAT and HESS at  $8..18 \text{ nW}/(\text{m}^2 \text{ srad})$   $\leftrightarrow$  tension with galaxy counts
- Indications ( $4 \sigma$ ) for excess transparency at large optical depth (both with Fermi-LAT and Cherenkov telescopes)
- Observations consistent with photon/ALPs mixing ( $m_a < 100 \text{ neV}$ ,  $g_{av} > 10^{-12} \text{ GeV}^{-1}$ )



# Outlook

- New VHE spectra available since ICRC 2013 (confirms and strengthens previous finding)
- Improved Fermi-LAT calibration/background rejection, more exposure, higher energy reach (up to 3 TeV) soon
- HESS-II accumulates data
- CTA in preparation (>2017)
- ALPS-II ideal laboratory experiment to search for gamma-ALPs mixing

