



# Holography with Large $N=4$

David Tong

Based on arXiv:1402.5135

# Holography with *Small* $N=4$

D1/D5 System



Near horizon limit:

$$AdS_3 \times S^3 \times \mathcal{M}$$

$$\mathcal{M} = T^4 \text{ or } K3$$

$Q_5$  D5-branes: 012345

$Q_1$  D1-branes: 05

# Holography with *Small* N=4

$$AdS_3 \times S^3 \times \mathcal{M}$$

$$\mathcal{M} = T^4 \text{ or } K3$$

## Boundary CFT

The central charge is:  $c = 6Q_5Q_1$

The R-symmetry becomes a current algebra

$$SO(4) \cong SU(2)_L \times SU(2)_R$$

Boundary theory has small N=4 superalgebra and is well understood

# Holography with *Large* $N=4$

$$AdS_3 \times S^3_+ \times S^3_- \times S^1$$

Supported by fluxes  $Q_5^\pm$  and  $Q_1$

Elitzur, Feinerman, Giveon and Tsabar (1998)  
de Boer and Skenderis (1999)  
Gukov, Martinec, Moore and Strominger (2004)

# Holography with *Large N=4*

The R-symmetry is now:

$$SO(4)^- \times SO(4)^+ \cong SU(2)_L^- \times SU(2)_R^- \times SU(2)_L^+ \times SU(2)_R^+$$

- There are two  $SU(2)$  current algebras.
- There is also a  $U(1)$  coming from  $S^1$  factor of the geometry.

Boundary theory has large  $N=4$  superalgebra. But what is the theory?!

Elitzur, Feinerman, Giveon and Tsabar (1998)  
de Boer and Skenderis (1999)  
Gukov, Martinec, Moore and Strominger (2004)

## Some Strange Properties of Large $N=4$

# Central Charge

$AdS_3 \times S^3_+ \times S^3_- \times S^1$  supported by fluxes  $Q_5^\pm$  and  $Q_1$

$$c = 6Q_1 \frac{Q_5^+ Q_5^-}{Q_5^+ + Q_5^-}$$

# BPS Bound

$AdS_3 \times S^3_+ \times S^3_- \times S^1$  supported by fluxes  $Q_5^\pm$  and  $Q_1$



$$h = \frac{Q_5^-}{Q_5^- + Q_5^+} l^+ + \frac{Q_5^+}{Q_5^- + Q_5^+} l^- + \frac{1}{Q_1(Q_5^+ + Q_5^-)} [(l^+ - l^-)^2 + u^2]$$

Representation under  $SU(2)^{+/-}$ 
Charge under  $U(1)$

- The bound is non-linear.
  - There is no chiral ring!
- The non-linearities are “1/N” suppressed.
  - They are not seen in supergravity


# How to Build the Boundary Field Theory

# A Tantalising D-Brane Configuration



$Q_5^+$  D5-branes: 012345  
 $Q_5^-$  D5'-branes: 056789  
 $Q_1$  D1-branes: 05

# Taking the Near Horizon Limit



$Q_5^+$  D5-branes: 012345  
 $Q_5^-$  D5'-branes: 056789  
 $Q_1$  D1-branes: 05


Smear D1-branes along 1234 and 6789. The near horizon limit is:

$$AdS_3 \times \mathbf{S}_+^3 \times \mathbf{S}_-^3 \times \mathbf{R}$$

How should we interpret this?!

Basic Idea: Study this D-brane configuration anyway!

# The Low-Energy Dynamics of D-Branes




$Q_5^+$  D5-branes: 012345  
 $Q_5^-$  D5'-branes: 056789  
 $Q_1$  D1-branes: 05

D1-D1 strings:  $U(Q_1)$  vector multiplet

- $N=(8,8)$  supersymmetry
- Gauge field and four complex, adjoint scalars

# The Low-Energy Dynamics of D-Branes




$Q_5^+$  D5-branes: 012345  
 $Q_5^-$  D5'-branes: 056789  
 $Q_1$  D1-branes: 05

D1-D5 strings:  $Q_5^+$  fundamental hypermultiplets

- $N=(4,4)$  supersymmetry
- two complex fundamental scalars

# The Low-Energy Dynamics of D-Branes




$Q_5^+$  D5-branes: 012345  
 $Q_5^-$  D5'-branes: 056789  
 $Q_1$  D1-branes: 05

D1-D5' strings:  $Q_5^-$  fundamental (twisted) hypermultiplets

- $N=(4,4)$  supersymmetry (but a different one!)
- two complex fundamental scalars

# The Low-Energy Dynamics of D-Branes




$Q_5^+$  D5-branes: 012345  
 $Q_5^-$  D5'-branes: 056789  
 $Q_1$  D1-branes: 05

D5-D5' strings:  $Q_5^+ Q_5^-$  Fermi multiplets

- $N=(0,8)$  supersymmetry
- only left-moving fermions

# The Low-Energy Dynamics of D-Branes



- All branes together preserve  $N=(0,4)$  supersymmetry
- The only question: how do chiral fermions couple to the other fields?
  - Surprising answer: the coupling is fixed by supersymmetry

# $N=(0,4)$ $d=1+1$ $U(Q_1)$ Gauge Theory

- $Q_5^+$  fundamental hypermultiplets
- $Q_5^-$  fundamental twisted hypermultiplets
- $Q_5^+Q_5^-$  neutral Fermi multiplets

Flavour symmetry:  $SU(Q_5^+) \times SU(Q_5^-)$

Also:  $SO(4)^- \times SO(4)^+ \cong SU(2)_L^- \times SU(2)_R^- \times SU(2)_L^+ \times SU(2)_R^+$

$(Z, \tilde{Z})$  in  $(2,2)$



$(Y, \tilde{Y})$  in  $(2,2)$



These are  $N=(0,4)$  R-symmetries



And...the theory has a  $U(1)$  global flavour symmetry which rotates Fermi multiplets and other stuff

Note: This is a chiral gauge theory. Which is not what we're looking for!


Flowing to the Infra-Red

# Computing the Central Charge

- Gauge theories in  $d=1+1$  are not conformal. But they can flow to conformal field theories.
- It's possible to compute the central charge for supersymmetric gauge theories. (At least  $N=(0,2)$ )
- The OPE of the right-moving R-current includes the term

$$R(x)R(y) \sim \frac{3c_R}{(x^- - y^-)^2} + \dots$$

But this is the anomaly. Which means that the central charge can be computed in the ultra-violet

$$c_R = 3\text{Tr } R^2$$


Sum over right-moving fermions, minus left-moving fermions

# Computing the Central Charge of $N=(0,2)$ Theories


$$c_R = 3\text{Tr } R^2$$

There's one small catch: you have to identify the right R-current in the UV. The requirement is:

$$\text{Tr } FR = 0$$

Silverstein and Witten (1993)

Any  $U(1)$  global symmetry  $F$



This can be repackaged as “c-extremization”

Adams, Tong and Wecht  
Benini and Bobev (2012)

Subtlety: In  $N=(0,4)$  theories, the R-symmetry is non-abelian.

# Computing the Central Charge of our $N=(0,4)$ Theory

$$c_R = 3\text{Tr } R^2 \quad \Rightarrow$$

$$c = 6Q_1 \frac{Q_5^+ Q_5^-}{Q_5^+ + Q_5^-}$$

In agreement with the result for a large  $N=(4,4)$  theory.  
(And in agreement with supergravity)

# Other Aspects of the $N=(0,4)$ Theory

## Things that work

Our theory has an  $SO(4) \times SO(4)$  symmetry.

The anomaly in UV symmetry should coincide with level of current algebra at fixed point.

$$SU(2)_R^+ \text{ and } SU(2)_L^+ : \quad \text{Tr} (R^+)^2 = \text{Tr} (L^+)^2 = Q_1 Q_5^+$$

$$SU(2)_R^- \text{ and } SU(2)_L^- : \quad \text{Tr} (R^-)^2 = \text{Tr} (L^-)^2 = Q_1 Q_5^-$$


## Things that don't immediately work

- There is no symmetry corresponding to the  $S^1$  action of the geometry.
- The  $N=(0,4)$  theory is chiral. It has a left-moving central charge given by

$$c_L = c_R + 2Q_5^+ Q_5^-$$


# Proposal

$N=(0,4)$  Gauge Theory



Large  $N=4$  CFT dual to  $\text{AdS}_3 \times S^3 \times S^3 \times S^1$   
+ decoupled left-moving fermions

# Where does this CFT live?




Can show that the CFT degrees of freedom are localised at the origin.

- This is slightly surprising in  $d=1+1$  (Mermin-Wagner theorem)
- But 5-branes come with infinitely long throats and related things are known to happen in  $N=(4,4)$ 
  - Decoupling of Higgs and Coulomb branches

# Work in Progress

Work with Ofer Aharony and Kenny Wong



We want to understand the dynamics of D-strings at the intersection.

Close to the D5-branes, the D-string feels an infinitely long throat. This is the region where throats meet.

# Work in Progress

Work with Ofer Aharony and Kenny Wong

The  $U(1)$  gauge theory on a single D-string is more tractable.

To understand the throat region, work with the Coulomb branch variables and integrate out charged fields

$$\begin{aligned} ds^2 &= \left( \frac{1}{g^2} + \frac{Q_5^+}{y^2} \right) (dy^2 + y^2 (d\Omega_3^+)^2) + \left( \frac{1}{g^2} + \frac{Q_5^-}{z^2} \right) (dz^2 + z^2 (d\Omega_3^-)^2) \\ &\longrightarrow Q_5^+ \left( \frac{dy^2}{y^2} + (d\Omega_3^+)^2 \right) + Q_5^- \left( \frac{dz^2}{z^2} + (d\Omega_3^-)^2 \right) \end{aligned}$$

There are also torsion terms and background dilaton charge.

The chiral fermions couple the two throats together. In the limit where the two throats meet, we have:

- The right number of chiral fermions decouples from the dynamics!
- Things are strongly coupled
- We still don't know how to see the extra  $S^1$  symmetry

# Summary and Open Problems

$N=(0,4)$  Gauge Theory



Large  $N=4$  CFT dual to  $\text{AdS}_3 \times S^3 \times S^3 \times S^1$   
+ decoupled left-moving fermions

- Central charge and anomalies all correct
- Good evidence for decoupling of chiral sector
- But no idea how the extra  $U(1)$  current may emerge

Future Questions:

- Compare chiral primaries
  - Marginal Operators
  - Any sign of non-linear BPS bound?
- Regimes of parameters where  $c$  scales linearly.
  - Relationship to higher spin theories?
- More complication  $\text{AdS}_3$  string solutions with  $N=(4,2)$ 
  - Related  $N=(0,2)$  gauge theories?

$$c = 6Q_1 \frac{Q_5^+ Q_5^-}{Q_5^+ + Q_5^-}$$

Gaberdiel and Gopakumar (2013)

Donos and Gauntlett (2013)

Thank you for your attention

# The Algebra

$$G^a(z)G^b(w) = \frac{2c}{3} \frac{\delta^{ab}}{(z-w)^3} - \frac{8\gamma\alpha_{ab}^{+,i}A^{+,i}(w) + 8(1-\gamma)\alpha_{ab}^{-,i}A^{-,i}(w)}{(z-w)^2} - \frac{4\gamma\alpha_{ab}^{+,i}\partial A^{+,i}(w) + 4(1-\gamma)\alpha_{ab}^{-,i}\partial A^{-,i}(w)}{z-w} + \frac{2\delta^{ab}L(w)}{z-w} + \dots,$$

$$A^{\pm,i}(z)A^{\pm,j}(w) = -\frac{k^{\pm}\delta^{ij}}{2(z-w)^2} + \frac{\epsilon^{ijk}A^{\pm,k}(w)}{z-w} + \dots,$$

$$Q^a(z)Q^b(w) = -\frac{(k^+ + k^-)\delta^{ab}}{2(z-w)} + \dots,$$

$$U(z)U(w) = -\frac{k^+ + k^-}{2(z-w)^2} + \dots,$$

$$A^{\pm,i}(z)G^a(w) = \mp \frac{2k^{\pm}\alpha_{ab}^{\pm,i}Q^b(w)}{(k^+ + k^-)(z-w)^2} + \frac{\alpha_{ab}^{\pm,i}G^b(w)}{z-w} + \dots,$$

$$A^{\pm,i}(z)Q^a(w) = \frac{\alpha_{ab}^{\pm,i}Q^b(w)}{z-w} + \dots,$$

$$Q^a(z)G^b(w) = \frac{2\alpha_{ab}^{+,i}A^{+,i}(w) - 2\alpha_{ab}^{-,i}A^{-,i}(w)}{z-w} + \frac{\delta^{ab}U(w)}{z-w} + \dots,$$

$$U(z)G^a(w) = \frac{Q^a(w)}{(z-w)^2} + \dots$$

# N=(0,2) Supersymmetry

Use N=(0,2) superfields. Scalar potential terms are built using Fermi multiplets.

$$\Psi = \psi_- - \theta^+ G - i\theta^+ \bar{\theta}^+ (D_0 + D_1)\psi_- - \bar{\theta}^+ E(\phi_i) + \theta^+ \bar{\theta}^+ \frac{\partial E}{\partial \phi^i} \psi_{+i}$$

E-Terms

$$\bar{D}_+ \Psi_a = E_a(\Phi_i)$$

Holomorphic functions of chiral superfields

J-Terms

$$S_J = \int d^2x d\theta^+ \sum_a \Psi_a J^a(\Phi_i) + \text{h.c.}$$

The scalar potential is:

$$V = \sum_a |E_a|^2 + |J_a|^2 + D^2$$

But there is only N=(0,2) supersymmetry if:

$$E \cdot J \equiv \sum_a E_a J^a = 0.$$

# The Coupling of the Chiral Fermions

The D5-D5' strings are needed to ensure that  $E.J=0$

$$E_{\chi} = -\frac{1}{2}\tilde{\Phi}\Phi'$$

$$J_{\chi} = \frac{1}{2}\tilde{\Phi}'\Phi$$

$$E_{\tilde{\chi}} = -\frac{1}{2}\tilde{\Phi}'\Phi$$

$$J_{\tilde{\chi}} = \frac{1}{2}\tilde{\Phi}\Phi'$$

# The Scalar Potential

To write it in a way in which the symmetries are manifest, define:

$$\omega = \begin{pmatrix} \phi \\ \tilde{\phi}^\dagger \end{pmatrix} \quad \omega' = \begin{pmatrix} \phi' \\ \tilde{\phi}'^\dagger \end{pmatrix}$$

and

$$\vec{D}_Z = \vec{\eta}_{ij} Z^i Z^j + \omega^\dagger \vec{\sigma} \omega$$

$\nearrow$   
 In  $\mathbf{3}$  of  $SU(2)_R^+$

$$\vec{D}_Y = \vec{\eta}_{ij} Y^i Y^j + \omega'^\dagger \vec{\sigma} \omega'$$


$\nearrow$        $\nwarrow$        $\nwarrow$   
 In  $\mathbf{3}$  of  $SU(2)_R^-$       Self-dual 't Hooft matrices      Pauli matrices

$$V = \text{Tr} (\vec{D}_Z \cdot \vec{D}_Z + \vec{D}_Y \cdot \vec{D}_Y) + \omega^\dagger Y^i Y^i \omega + \omega'^\dagger Z^i Z^i \omega' + \text{Tr} [Y^i, Z^j]^2 + \text{Tr} (\omega^\dagger \cdot \omega \omega'^\dagger \cdot \omega')$$


$$\text{last term} = \sum_{a=1}^{Q_5^+} \sum_{b=1}^{Q_5^-} \left( (\phi_a^\dagger \phi_b') (\phi_b'^\dagger \phi_a) + (\tilde{\phi}_a \tilde{\phi}_b'^\dagger) (\tilde{\phi}_b' \tilde{\phi}_a^\dagger) + (\phi_a^\dagger \tilde{\phi}_b'^\dagger) (\tilde{\phi}_b' \phi_a) + (\tilde{\phi}_a \phi_b') (\phi_b'^\dagger \tilde{\phi}_a^\dagger) \right)$$

# Vacuum Moduli Space


$$V = \text{Tr} (\vec{D}_Z \cdot \vec{D}_Z + \vec{D}_Y \cdot \vec{D}_Y) + \omega^\dagger Y^i Y^i \omega + \omega'^\dagger Z^i Z^i \omega' + \text{Tr} [Y^i, Z^j]^2 + \text{Tr} (\omega^\dagger \cdot \omega \omega'^\dagger \cdot \omega')$$



- $\omega = \omega' = 0$  with  $Z^i$  and  $Y^i$  mutually commuting.



- $\vec{D}_Z = 0$  and  $Y^i = \omega' = 0$



- $\vec{D}_Y = 0$  and  $Z^i = \omega = 0$

But in fact, we'll be interested in modes which appear to localise at the origin...

# Computing the Central Charge of $N=(0,4)$ Theories

Right-moving R-current is:  $SU(2)_R^- \times SU(2)_R^+$

- No mixing for non-Abelian symmetries
- But  $N=(0,2)$  R-current is some combination of

$$R^\pm \subset su(2)_R^\pm$$

Both are good  $N=(0,2)$  R-currents. But there is one combination that is an  $N=(0,2)$  flavour symmetry

$$U = R^+ - R^-$$


We must have

$$\text{Tr } UR = 0$$

$$\begin{array}{l} \text{Tr } UR^- = -2Q_1 Q_5^- \\ \text{Tr } UR^+ = +2Q_1 Q_5^+ \end{array} \quad \Rightarrow \quad R = \frac{Q_5^+}{Q_5^+ + Q_5^-} R^- + \frac{Q_5^-}{Q_5^+ + Q_5^-} R^+$$

# What happened to the Chiral Modes?

An interesting story was found in the absence of D1-branes. This configuration is called the *I-brane*



- Chiral modes are pushed away from the intersection.
- The intersection has a mass gap
- It also has  $d=2+l$  dimensional symmetry!