

# Diffusion and Chaos in Quantum Matter

Mike Blake - MIT



MAB -1603.08510 (PRL)

MAB -1604.01754

MAB and Donos - 1611.09380

MAB, Davison and Sachdev -1705.07896

# Motivation

- Evidence has recently emerged for a new connection between transport (conductivities etc) and many-body chaos.
- Long-standing idea that dynamics at strong coupling governed by 'Planckian' mean free time

$$\tau \sim \frac{\hbar}{k_B T}$$

- In a Fermi liquid thermoelectric diffusion constants are of form

$$D \sim v_F^2 \tau_{FL}$$

- **Hartnoll** (2014) proposed that strange metal transport is governed by

$$D \sim \frac{\hbar v^2}{k_B T}$$

- But unclear how to identify characteristic velocity of a strongly coupled theory.

# Proposal

- A natural infra-red velocity is provided by speed at which chaos propagates

$$\langle [\hat{W}_x(t_w), \hat{V}_y(0)]^2 \rangle_\beta \sim f_1 e^{\lambda_L(t_w - t_* - |x-y|/v_B)}$$

- Suggest butterfly velocity as characteristic velocity for diffusion

$$D \sim v_B^2 \tau$$


- In many cases also expect Lyapunov exponent to provide relevant mean free time  $\tau \sim \tau_L = \lambda_L^{-1}$

# Holography

Classical gravity in  
asymptotically-AdS spacetime



Strongly coupled  
large N gauge theory



- Reissner-Nordstrom black hole
- Hawking radiation = finite temperature,  $T$
- Electric field = chemical potential,  $\mu$

- DC thermoelectric conductivities  $\sigma \propto \bar{\kappa}$  can be related to geometry and fields at black hole horizon.

MAB & Tong;  
Donos & Gauntlett

- Likewise chaos exponents can be calculated from gravitational shock-wave on horizon.

Shenker & Stanford;  
Roberts, Stanford &  
Susskind

- Diffusion constants proportional to conductivities through Einstein relations.

# Holographic examples

- First evidence came from charge diffusion in particle-hole symmetric theories.

$$D_c = \frac{\sigma}{\chi} \quad \chi = \left( \frac{\partial \rho}{\partial \mu} \right)_T$$

- Calculated this for holographic theories that flow to Lifshitz/hyperscaling geometries in IR.
- Generalised scaling theories described by critical exponents  $(z, \theta, \phi)$

- Shock-wave calculation for these geometries gives

$$\tau_L^{-1} = 2\pi T \quad v_B^2 \sim L^2 T^{2-2/z}$$

- Found these chaos parameters were universally related to the diffusion constant

$$D_c = \frac{d_\theta}{\Delta_\chi} v_B^2 \tau_L$$

MAB

- Although such a simple relationship is quite special to particle-hole symmetric theories.



- More general connection is found in relationship between energy/thermal diffusion and chaos

$$D_T = \frac{\kappa}{c_\rho} \quad c_\rho = T \left( \frac{\partial s}{\partial T} \right)_\rho$$

- For these scaling geometries this is always given by

$$D_T = \frac{z}{2z - 2} v_B^2 \tau_L$$

MAB;  
MAB, Davison and Sachdev


- This holds independently of charge density, magnetic field, periodic potential strength.

Fixed points with  $z=1$  (e.g. CFTs) are a special exception

# Other Examples

## SYK models

Sachdev & Ye;  
Kitaev;  
Stanford & Maldacena...



$$H = \sum_{1 \leq j < k < l < m \leq N} J_{jklm} \chi_j \chi_k \chi_l \chi_m$$

$$\frac{1}{3!} N^3 \overline{J_{jklm}^2} = J^2$$

- Infra-red physics dominated by 'Schwarzian' mode


$$t \rightarrow t + \epsilon(t) \quad \tau_L^{-1} = 2\pi T$$

- Same mode and action are found in near-AdS2 black holes.

Jensen;  
Stanford, Maldacena & Yang

- Higher dimensional generalisation consists of a chain of coupled SYK models

Gu, Stanford & Qi;  
Davison et al



- Schwarzian mode  $t \rightarrow t + \epsilon_x(t)$  gives both chaos exponents and thermal diffusivity.

$$\tau_L^{-1} = 2\pi T \quad v_B^2 \sim \frac{J'^2 T}{J} \quad D_T = v_B^2 \tau_L$$

- Perfect match with results for holographic theories that flow to  $AdS_2 \times R^{d-1}$

# Critical Fermi surfaces

Patel & Sachdev

## Quantum chaos on a critical Fermi surface

Aavishkar A. Patel<sup>1</sup> and Subir Sachdev<sup>1,2</sup>

<sup>1</sup>*Department of Physics, Harvard University, Cambridge MA 02138, USA*

<sup>2</sup>*Perimeter Institute for Theoretical Physics,  
Waterloo, Ontario, Canada N2L 2Y5*

### Abstract

We compute parameters characterizing many-body quantum chaos for a critical Fermi surface without quasiparticle excitations. We examine a theory of  $N$  species of fermions at non-zero density coupled to a  $U(1)$  gauge field in two spatial dimensions, and determine the Lyapunov rate and the butterfly velocity in an extended random-phase approximation. The thermal diffusivity is found to be universally related to these chaos parameters *i.e.* the relationship is independent of  $N$ , the gauge coupling constant, the Fermi velocity, the Fermi surface curvature, and high energy details.

$$\mathcal{L}[\Psi, a] = \Psi^\dagger \left( \partial_\tau - ia_\tau - \frac{(\nabla - i\vec{a})^2}{2m} - \mu \right) \Psi + \frac{1}{2g^2} (\nabla \times \vec{a})^2$$

$$\tau_L = \frac{\hbar}{2.48 k_B T}$$

$$v_B \sim \frac{N v_F^{5/3}}{e^{4/3} \gamma^{1/3}} T^{1/3}$$

$$D_E = 0.42 v_B^2 \tau_L$$

str-el] 12 Mar 2017

- Connections of the form  $D \sim v_B^2 \tau_L$  have also been seen in

Electron/phonon bad metals - Werman, Kivelson & Berg

Incoherent Bose-Hubbard models - Bohrdt, Endrel, Mendes & Knap

O(N) models - Chowdury & Swingle

Fermi liquids with disorder/electron-electron interactions - Aleiner, Faoro & Ioffe

Higher derivative holographic theories - Baggioli, Gouteraux, Kiritsis & Li

- Recent work provides evidence for a relationship between a light cone velocity and diffusion.

Hartman, Hartnoll & Mahajan  
Lucas

**Thank you!**