

# Hyperbolic Extra Dimensions at the Large Hadron Collider

Henrik Melb us  
melbeus@kth.se

Royal Institute of Technology (KTH), Stockholm, Sweden

Nordic Workshop on LHC and Beyond  
June 12, 2008

# Outline

## Outline

- Motivation and description of the model
- Possible signals at the LHC
- Conclusions

## The paper

- Based on [arXiv:0806.1841 \[hep-ph\]](#)
- In collaboration with Tommy Ohlsson

# The hierarchy problem

- The electroweak scale  $M_{\text{ew}} \sim 100 \text{ GeV}$
- The Planck scale  $M_{\text{Pl}} \equiv G_N^{-1/2} \sim 10^{18} \text{ GeV}$

# The hierarchy problem

- The electroweak scale  $M_{\text{ew}} \sim 100 \text{ GeV}$
- The Planck scale  $M_{\text{Pl}} \equiv G_N^{-1/2} \sim 10^{18} \text{ GeV}$
- Theoretically,  $M_{\text{ew}}$  is expected to obtain loop corrections of order  $M_{\text{Pl}} \gg M_{\text{ew}}$
- Miraculous cancellation needed to keep  $M_{\text{ew}}$  at  $\sim 100 \text{ GeV}$

# The hierarchy problem

- The electroweak scale  $M_{\text{ew}} \sim 100 \text{ GeV}$
- The Planck scale  $M_{\text{Pl}} \equiv G_N^{-1/2} \sim 10^{18} \text{ GeV}$
- Theoretically,  $M_{\text{ew}}$  is expected to obtain loop corrections of order  $M_{\text{Pl}} \gg M_{\text{ew}}$
- Miraculous cancellation needed to keep  $M_{\text{ew}}$  at  $\sim 100 \text{ GeV}$
- Popular solution: SUSY (cancellation of divergences)

# The hierarchy problem

- The electroweak scale  $M_{\text{ew}} \sim 100 \text{ GeV}$
- The Planck scale  $M_{\text{Pl}} \equiv G_N^{-1/2} \sim 10^{18} \text{ GeV}$
- Theoretically,  $M_{\text{ew}}$  is expected to obtain loop corrections of order  $M_{\text{Pl}} \gg M_{\text{ew}}$
- Miraculous cancellation needed to keep  $M_{\text{ew}}$  at  $\sim 100 \text{ GeV}$
- Popular solution: SUSY (cancellation of divergences)
- Alternative solution: **Large extra dimensions**

# Extra dimensions — the ADD model

- The ADD model:
  - ▶ Arkani-Hamed, Dimopolous, and Dvali (1998)
  - ▶  $d$  extra spatial dimensions
  - ▶ Compactified on a torus (a flat space)
  - ▶ SM fields confined to a 4-dimensional brane
  - ▶ Only gravity can probe the extra dimensions

# Extra dimensions — the ADD model

- The ADD model:
  - ▶ Arkani-Hamed, Dimopolous, and Dvali (1998)
  - ▶  $d$  extra spatial dimensions
  - ▶ Compactified on a torus (a flat space)
  - ▶ SM fields confined to a 4-dimensional brane
  - ▶ Only gravity can probe the extra dimensions
- The extra dimensions can be large,  $\sim 10 \mu\text{m}$ , i.e.,  
Large extra dimensions

# Solution to the hierarchy problem

- Because gravity propagates in  $4 + d$  dimensions, the Planck scale is only an effective scale valid at low energies
- New fundamental mass scale for gravity,  $M_*$

# Solution to the hierarchy problem

- Because gravity propagates in  $4 + d$  dimensions, the Planck scale is only an effective scale valid at low energies
- New fundamental mass scale for gravity,  $M_*$
- $M_{\text{Pl}}^2 = VM_*^{2+d} = (2\pi L)^d M_*^{2+d}$
- If  $V$  is sufficiently large, then we could have  $M_* \sim M_{\text{ew}}$

# Solution to the hierarchy problem

- Because gravity propagates in  $4 + d$  dimensions, the Planck scale is only an effective scale valid at low energies
- New fundamental mass scale for gravity,  $M_*$
- $M_{\text{Pl}}^2 = VM_*^{2+d} = (2\pi L)^d M_*^{2+d}$
- If  $V$  is sufficiently large, then we could have  $M_* \sim M_{\text{ew}}$
- Large extra dimensions expected at  $\sim 1$  TeV (like SUSY)

# Solution to the hierarchy problem

- Because gravity propagates in  $4 + d$  dimensions, the Planck scale is only an effective scale valid at low energies
- New fundamental mass scale for gravity,  $M_*$
- $M_{\text{Pl}}^2 = VM_*^{2+d} = (2\pi L)^d M_*^{2+d}$
- If  $V$  is sufficiently large, then we could have  $M_* \sim M_{\text{ew}}$
- Large extra dimensions expected at  $\sim 1$  TeV (like SUSY)
- **However**, if  $M_* = 1$  TeV, then  $L \sim 10^{31/d} \text{ TeV}^{-1} \gg M_*^{-1}$
- **New hierarchy** between  $M_*$  and  $L^{-1}$

# Hyperbolic extra dimensions

- Hyperbolic geometry: constant negative curvature  $v$
- Hyperbolic disc of radius  $L$ :

$$V = \frac{4\pi}{v^2} \sinh^2 \left( \frac{vL}{2} \right) \sim \frac{\pi}{v^2} \exp(vL)$$

- Possible to have  $M_* \sim M_{\text{ew}}$  and  $M_* \sim L^{-1}$

# Hyperbolic extra dimensions

- Hyperbolic geometry: constant negative curvature  $v$
- Hyperbolic disc of radius  $L$ :

$$V = \frac{4\pi}{v^2} \sinh^2 \left( \frac{vL}{2} \right) \sim \frac{\pi}{v^2} \exp(vL)$$

- Possible to have  $M_* \sim M_{\text{ew}}$  and  $M_* \sim L^{-1}$
- Hyperbolic extra dimensions provide a more satisfactory solution to the hierarchy problem
  - ▶ Kaloper, March-Russell, Starkman, and Trodden (2000)

# Hyperbolic extra dimensions

- Hyperbolic geometry: constant negative curvature  $v$
- Hyperbolic disc of radius  $L$ :

$$V = \frac{4\pi}{v^2} \sinh^2 \left( \frac{vL}{2} \right) \sim \frac{\pi}{v^2} \exp(vL)$$

- Possible to have  $M_* \sim M_{\text{ew}}$  and  $M_* \sim L^{-1}$
- Hyperbolic extra dimensions provide a more satisfactory solution to the hierarchy problem
  - ▶ Kaloper, March-Russell, Starkman, and Trodden (2000)
- Our model: two extra dimensions with the geometry of a **hyperbolic disc**

# Comparison to the ADD model

## The ADD model

- 1 free parameter:  $M_*$

## The hyperbolic disc model

- 3 free parameters:  $M_*$ ,  $\nu$ , and  $\tau_b$

# Comparison to the ADD model

## The ADD model

- 1 free parameter:  $M_*$
- KK spectrum starts out at  $m \approx 0$

## The hyperbolic disc model

- 3 free parameters:  $M_*$ ,  $\nu$ , and  $\tau_b$
- KK spectrum starts out at  $m \approx \nu/2 > 0$

# Comparison to the ADD model

## The ADD model

- 1 free parameter:  $M_*$
- KK spectrum starts out at  $m \approx 0$
- For  $d = 2$ ,  $M_*$  is constrained to  $M_* \geq 50$  TeV from astrophysics

## The hyperbolic disc model

- 3 free parameters:  $M_*$ ,  $v$ , and  $\tau_b$
- KK spectrum starts out at  $m \approx v/2 > 0$
- Weak restrictions on the parameter space

# Comparison to the ADD model

## The ADD model

- 1 free parameter:  $M_*$
- KK spectrum starts out at  $m \approx 0$
- For  $d = 2$ ,  $M_*$  is constrained to  $M_* \geq 50$  TeV from astrophysics
- Universal coupling of KK modes

## The hyperbolic disc model

- 3 free parameters:  $M_*$ ,  $\nu$ , and  $\tau_b$
- KK spectrum starts out at  $m \approx \nu/2 > 0$
- Weak restrictions on the parameter space
- Different KK modes have different couplings to SM fields

# Comparison to the ADD model

## The ADD model

- 1 free parameter:  $M_*$
- KK spectrum starts out at  $m \approx 0$
- For  $d = 2$ ,  $M_*$  is constrained to  $M_* \geq 50$  TeV from astrophysics
- Universal coupling of KK modes
- Physical results independent of the position of the brane

## The hyperbolic disc model

- 3 free parameters:  $M_*$ ,  $v$ , and  $\tau_b$
- KK spectrum starts out at  $m \approx v/2 > 0$
- Weak restrictions on the parameter space
- Different KK modes have different couplings to SM fields
- Physical results depend on the position of the brane

# LHC phenomenology

- The LHC phenomenology for the ADD model has been investigated
  - ▶ Giudice, Rattazzi, and Wells (2000)
  - ▶ Hinchliffe and Vacavant (2001)

# LHC phenomenology

- The LHC phenomenology for the ADD model has been investigated
  - ▶ Giudice, Rattazzi, and Wells (2000)
  - ▶ Hinchliffe and Vacavant (2001)
- Interesting reactions at the LHC:
  - ▶  $p + p \rightarrow \text{jet} + G$
  - ▶  $p + p \rightarrow \gamma + G$   
where  $G$  is a graviton
- The graviton is not observed — missing energy signatures


# LHC phenomenology

- The LHC phenomenology for the ADD model has been investigated
  - ▶ Giudice, Rattazzi, and Wells (2000)
  - ▶ Hinchliffe and Vacavant (2001)
- Interesting reactions at the LHC:
  - ▶  $p + p \rightarrow \text{jet} + G$
  - ▶  $p + p \rightarrow \gamma + G$   
where  $G$  is a graviton
- The graviton is not observed — missing energy signatures
- We have studied the LHC phenomenology for the hyperbolic disc model

# LHC phenomenology


- The LHC phenomenology for the ADD model has been investigated
  - ▶ Giudice, Rattazzi, and Wells (2000)
  - ▶ Hinchliffe and Vacavant (2001)
- Interesting reactions at the LHC:
  - ▶  $p + p \rightarrow \text{jet} + G$
  - ▶  $p + p \rightarrow \gamma + G$   
where  $G$  is a graviton
- The graviton is not observed — **missing energy** signatures
- We have studied the LHC phenomenology for the hyperbolic disc model
- The main SM background, from  $p + p \rightarrow \text{jet}/\gamma + Z(\rightarrow \nu\bar{\nu})$ , has been simulated in PYTHIA

# Cross sections — hadronic jet production




**Figure:** The differential cross section for graviton plus jet production with respect to  $p_T$ . Left panel:  $M_* = 1.5$  TeV. Right panel:  $M_* = 2$  TeV.

# Cross sections — hadronic jet production




**Figure:** The differential cross section for graviton plus jet production with respect to  $\cos(\theta)$ . Left panel:  $M_* = 1.5$  TeV. Right panel:  $M_* = 2$  TeV.

# Cross sections — photon production



**Figure:** The differential cross section for graviton plus photon production with respect to  $p_T$ . Left panel:  $M_* = 1$  TeV. Right panel:  $M_* = 1.5$  TeV.

## Cross sections — photon production



**Figure:** The differential cross section for graviton plus photon production with respect to  $\cos(\theta)$ . Left panel:  $M_* = 1$  TeV. Right panel:  $M_* = 1.5$  TeV.

# Summary and conclusions

- The ADD model provides a possible solution to the **hierarchy problem**
- Hyperbolic extra dimensions improve this solution

# Summary and conclusions

- The ADD model provides a possible solution to the **hierarchy problem**
- Hyperbolic extra dimensions improve this solution
- Large extra dimensions can be studied at the LHC
  - ▶ Signals similar to the ADD model
  - ▶ In some cases indistinguishable
  - ▶ Most promising signal from the **jet channel**
  - ▶ More difficult for the photon channel

# Summary and conclusions

- The ADD model provides a possible solution to the **hierarchy problem**
- Hyperbolic extra dimensions improve this solution
- Large extra dimensions can be studied at the LHC
  - ▶ Signals similar to the ADD model
  - ▶ In some cases indistinguishable
  - ▶ Most promising signal from the **jet channel**
  - ▶ More difficult for the photon channel
- Paper on **arXiv:0806.1841 [hep-ph]**