


General Dark Matter Electron Interactions in Graphene

**R. Catena, T. Emken, M. Matas, N. Spaldin, EU:
2303.15497 & 2303.15509**

Einar Urdshals, 16. June 2023

General Idea



Motivation

- Electron recoils are sensitive to dark matter (DM) masses $\gtrsim 1 \text{ MeV}$
- Need to discriminate between DM events and background
- Graphene-like targets can produce a strong daily modulation, a smoking gun signal of dark matter


DM Induced Electron Ejections from Graphene

- $R \propto \int f_\chi(\mathbf{v}, t) R_{\text{free}}(\mathbf{v}, \mathbf{q}, \mathbf{k}') \left| \psi(\ell, E_e) \right|^2$ R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15497
- $f_\chi(\mathbf{v}, t)$ is the DM velocity distribution
- R_{free} is the free particle response function and depends on the physics of free particles
- $\left| \psi(\ell, E_e) \right|^2$ contains all the material physics

DM Velocity Distribution

- Seen from the lab on Earth the DM velocity distribution is anisotropic and time dependent

- $f_\chi(\mathbf{v}, t) \propto \exp\left[-\frac{(\mathbf{v} + \mathbf{v}_e(t))^2}{v_0^2}\right] \times \Theta(v_{\text{esc}} - |\mathbf{v} + \mathbf{v}_e(t)|)$



Free Particle Response Function

- R_{free} is built from non-relativistic effective operators, allows covering arbitrary spin 0 and 1/2 DM models.
- Depends on the physics of the free particles, i.e. the DM particle and the final state electron

$$\mathcal{O}_1 = \mathbb{1}_{\chi e}$$

$$\mathcal{O}_3 = i\mathbf{S}_e \cdot \left(\frac{\mathbf{q}}{m_e} \times \mathbf{v}_{\text{el}}^\perp \right)$$

$$\mathcal{O}_4 = \mathbf{S}_\chi \cdot \mathbf{S}_e$$

$$\mathcal{O}_5 = i\mathbf{S}_\chi \cdot \left(\frac{\mathbf{q}}{m_e} \times \mathbf{v}_{\text{el}}^\perp \right)$$

$$\mathcal{O}_6 = \left(\mathbf{S}_\chi \cdot \frac{\mathbf{q}}{m_e} \right) \left(\mathbf{S}_e \cdot \frac{\mathbf{q}}{m_e} \right)$$

$$\mathcal{O}_7 = \mathbf{S}_e \cdot \mathbf{v}_{\text{el}}^\perp$$

$$\mathcal{O}_8 = \mathbf{S}_\chi \cdot \mathbf{v}_{\text{el}}^\perp$$

$$\mathcal{O}_9 = i\mathbf{S}_\chi \cdot \left(\mathbf{S}_e \times \frac{\mathbf{q}}{m_e} \right)$$

$$\mathcal{O}_{10} = i\mathbf{S}_e \cdot \frac{\mathbf{q}}{m_e}$$

$$\mathcal{O}_{11} = i\mathbf{S}_\chi \cdot \frac{\mathbf{q}}{m_e}$$

$$\mathcal{O}_{12} = \mathbf{S}_\chi \cdot \left(\mathbf{S}_e \times \mathbf{v}_{\text{el}}^\perp \right)$$

$$\mathcal{O}_{13} = i \left(\mathbf{S}_\chi \cdot \mathbf{v}_{\text{el}}^\perp \right) \left(\mathbf{S}_e \cdot \frac{\mathbf{q}}{m_e} \right)$$

$$\mathcal{O}_{14} = i \left(\mathbf{S}_\chi \cdot \frac{\mathbf{q}}{m_e} \right) \left(\mathbf{S}_e \cdot \mathbf{v}_{\text{el}}^\perp \right)$$

$$\mathcal{O}_{15} = i\mathcal{O}_{11} \left[\left(\mathbf{S}_e \times \mathbf{v}_{\text{el}}^\perp \right) \cdot \frac{\mathbf{q}}{m_e} \right]$$

Modelling Graphene



Tight Binding Approximation Density Functional Theory

- Approximation based on analytic atomic wave functions
 - Not self consistent
 - Accurate near the nucleus
 - Computationally cheap
 - Semi-analytic, not a black box
- Works by finding the electron density that minimises the energy of the system
 - Self consistent from first principles
 - Can not treat the electron wave-function close to the nucleus
 - Computationally expensive

Momentum Distribution in Graphene

Tight Binding Approximation


Density Functional Theory



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15497

Total Rate of Ejected Electrons

Time Averaged Rate




\mathcal{O}_1 contact is dark photon with heavy mediator

\mathcal{O}_1 long range is dark photon with light mediator

$$\mathcal{O}_3 \propto \mathbf{q} \times \mathbf{v}$$

Daily Modulation



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15497

We recommend using
DFT, and all rates will from
now be DFT-obtained

Statistical Tests


Establishing Daily Modulation Excluding Parameter Space

- Divide the data set in two parts, n_+ and n_-
- No modulation
 $\implies E[n_+] = E[n_-]$
- Wish to reject the hypothesis that n_+ and n_- are drawn from the same distribution
- Exclude DM unlikely to produce $n_+ + n_-$ or fewer events

Specific Detector Setups

Detector Setups


Fixed Graphene



Fixed graphene sheets

Detector Setups

Moving Graphene



Moving graphene sheets


R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509

Detector Setups

Fixed Carbon NanoTubes (CNTs)


n_+ : Number of detected events
in 12 hours with most events

n_- : Number of detected events
in 12 hours with least events



Detector Setups


Moving Carbon NanoTubes (CNTs)




R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509

Potential for Establishing Modulation at 3σ

\mathcal{O}_1 Contact Interaction



\mathcal{O}_1 Long Range Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509


Corresponds to dark photon model
with a heavy mediator

10 g year exposure


Corresponds to dark photon model
with a light mediator

Potential for Establishing Modulation at 3σ

\mathcal{O}_3 Contact Interaction



Anapole Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509


$$\mathcal{O}_3 \propto \mathbf{q} \times \mathbf{v}$$

10 g year exposure


$$L_{\text{Anapole}} = \frac{g}{2\Lambda^2} \bar{\chi} \gamma^\mu \gamma^5 \chi \partial^\nu F_{\mu\nu}$$

Potential for Establishing Modulation at 3σ

Electric Dipole Interaction



Magnetic Dipole Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509


$$L_{\text{Electric dipole}} = \frac{g}{\Lambda} i \bar{\chi} \sigma^{\mu\nu} \gamma^5 \chi F_{\mu\nu}$$

10 g year exposure


$$L_{\text{Magnetic dipole}} = \frac{g}{\Lambda} \bar{\chi} \sigma^{\mu\nu} \chi F_{\mu\nu}$$

Exclusion Potential

\mathcal{O}_1 Contact Interaction



\mathcal{O}_1 Long Range Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509


Corresponds to dark photon model
with a heavy mediator

10 g year exposure


Corresponds to dark photon model
with a light mediator

Exclusion Potential

\mathcal{O}_3 Contact Interaction



Anapole Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509


$$\mathcal{O}_3 \propto \mathbf{q} \times \mathbf{v}$$

10 g year exposure


$$L_{\text{Anapole}} = \frac{g}{2\Lambda^2} \bar{\chi} \gamma^\mu \gamma^5 \chi \partial^\nu F_{\mu\nu}$$

Exclusion Potential

Electric Dipole Interaction



Magnetic Dipole Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509

$$L_{\text{Electric dipole}} = \frac{g}{\Lambda} i \bar{\chi} \sigma^{\mu\nu} \gamma^5 \chi F_{\mu\nu}$$

10 g year exposure

$$L_{\text{Magnetic dipole}} = \frac{g}{\Lambda} \bar{\chi} \sigma^{\mu\nu} \chi F_{\mu\nu}$$


Summary

- Experiments based on graphene and CNTs are suitable to produce a smoking gun signal of DM
- The relative performance of the experimental setups depends on the form of the DM electron interaction. Need to consider non-standard interactions.
- For more details, see our papers [2303.15497](#) & [2303.15509](#)


Backup Slides

Electron Density

Tight Binding Approximation




Density Functional Theory




R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15497

Modelling Graphene

Momentum Density



Band Structure




Daily Modulation in Total Ejection Rates

$$L_{\text{Anapole}} = \frac{g}{2\Lambda^2} \bar{\chi} \gamma^\mu \gamma^5 \chi \partial^\nu F_{\mu\nu}$$

$$L_{\text{Electric dipole}} = \frac{g}{\Lambda} i \bar{\chi} \sigma^{\mu\nu} \gamma^5 \chi F_{\mu\nu}$$


$$L_{\text{Magnetic dipole}} = \frac{g}{\Lambda} \bar{\chi} \sigma^{\mu\nu} \chi F_{\mu\nu}$$

$$\mathcal{O}_3 \propto \mathbf{q} \times \mathbf{v}$$




Daily Modulation in Observed Events

Fixed Graphene



Fixed CNTs



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509

$$L_{\text{Anapole}} = \frac{g}{2\Lambda^2} \bar{\chi} \gamma^\mu \gamma^5 \chi \partial^\nu F_{\mu\nu}$$




$$L_{\text{Electric dipole}} = \frac{g}{\Lambda} i \bar{\chi} \sigma^{\mu\nu} \gamma^5 \chi F_{\mu\nu}$$

$$L_{\text{Magnetic dipole}} = \frac{g}{\Lambda} \bar{\chi} \sigma^{\mu\nu} \chi F_{\mu\nu}$$


$$\mathcal{O}_3 \propto \mathbf{q} \times \mathbf{v}$$

Scattering Directions Fixed Sheets

$\mathcal{O}_3 \propto \mathbf{v} \times \mathbf{q}$, contact type interaction, $m_\chi = 5 \text{ MeV}$







R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509



Scattering Directions Fixed Sheets





Magnetic Dipole Type Interaction, $m_\chi = 5 \text{ MeV}$



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509

Scattering Directions Fixed CNTs





$\mathcal{O}_3 \propto \mathbf{v} \times \mathbf{q}$, contact type interaction, $m_\chi = 5 \text{ MeV}$



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509

Scattering Directions Fixed CNTs

Magnetic Dipole Type Interaction, $m_\chi = 5 \text{ MeV}$




More CNT Based Setups

Fixed CNTs Facing Away From DM Wind

n_+ : Number of detected events
in 12 hours with most events


n_- : Number of detected events
in 12 hours with least events



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509

More CNT Based Setups

Moving CNTs, Back to Back



More CNT Based Setups

Moving CNTs, 90° Relative Orientation

n_- : Number of detected events


upper detector

v_e

DM wind at **all times**

Detector

\vec{E}



Substrate


Detector

\vec{E}

n_+ : Number of detected events
in lower detector

More CNT Based Setups


Moving CNTs, 145° relative orientation




n_- : Number of detected events
upper detector

Potential for Establishing Modulation at 3σ

\mathcal{O}_1 Contact Interaction



\mathcal{O}_1 Long Range Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509


Corresponds to dark photon model
with a heavy mediator

10 g year exposure


Corresponds to dark photon model
with a light mediator

Potential for Establishing Modulation at 3σ

\mathcal{O}_3 Contact Interaction



Anapole Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509


$$\mathcal{O}_3 \propto \mathbf{q} \times \mathbf{v}$$

10 g year exposure


$$L_{\text{Anapole}} \in \frac{g}{2\Lambda^2} \bar{\chi} \gamma^\mu \gamma^5 \chi \partial^\nu F_{\mu\nu}$$

Potential for Establishing Modulation at 3σ

Electric Dipole Interaction



Magnetic Dipole Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509


$$L_{\text{Electric dipole}} \in \frac{g}{\Lambda} i \bar{\chi} \sigma^{\mu\nu} \gamma^5 \chi F_{\mu\nu}$$

10 g year exposure


$$L_{\text{Magnetic dipole}} \in \frac{g}{\Lambda} \bar{\chi} \sigma^{\mu\nu} \chi F_{\mu\nu}$$

Exclusion Potential

\mathcal{O}_1 Contact Interaction



\mathcal{O}_1 Long Range Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509


Corresponds to dark photon model
with a heavy mediator

10 g year exposure


Corresponds to dark photon model
with a light mediator

Exclusion Potential

\mathcal{O}_3 Contact Interaction



Anapole Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509


$$\mathcal{O}_3 \propto \mathbf{q} \times \mathbf{v}$$

10 g year exposure


$$L_{\text{Anapole}} \in \frac{g}{2\Lambda^2} \bar{\chi} \gamma^\mu \gamma^5 \chi \partial^\nu F_{\mu\nu}$$

Exclusion Potential

Electric Dipole Interaction



Magnetic Dipole Interaction



R. Catena, T. Emken, M. Matas, N. Spaldin, EU: 2303.15509

$$L_{\text{Electric dipole}} \in \frac{g}{\Lambda} i \bar{\chi} \sigma^{\mu\nu} \gamma^5 \chi F_{\mu\nu}$$

10 g year exposure

$$L_{\text{Magnetic dipole}} \in \frac{g}{\Lambda} \bar{\chi} \sigma^{\mu\nu} \chi F_{\mu\nu}$$