

Spectral Features From Pulsars and Dark Matter in the Local Cosmic-Ray Electron and Positron Flux

arXiv:2206.04699 & arXiv:2304.07317



Isabelle John
isabelle.john@fysik.su.se


With Tim Linden

17 October 2023
OKC@15 Stockholm




Cosmic-Ray Electrons and Positrons

Experimental




Cosmic-Ray Electrons and Positrons

Experimental





Modelling



Propagation and Energy Losses

Positron source
e.g. pulsars or
dark matter

Propagation and
Energy Losses



Synchrotron radiation
in magnetic fields


Inverse-Compton scattering
on ambient photons

Spectrum of an Individual Pulsar

1. Large fraction of positrons is produced when pulsar is very young



Spectrum of an Individual Pulsar

1. Large fraction of positrons is produced when pulsar is very young
2. High-energy positrons lose energy faster than low-energy positrons





Spectrum of an Individual Pulsar

1. Large fraction of positrons is produced when pulsar is very young
2. High-energy positrons lose energy faster than low-energy positrons
3. These initial positrons build up sharp feature in spectrum over time





Spectrum of an Individual Pulsar

1. Large fraction of positrons is produced when pulsar is very young
2. High-energy positrons lose energy faster than low-energy positrons
3. These initial positrons build up sharp feature in spectrum over time





Spectrum of an Individual Pulsar

1. Large fraction of positrons is produced when pulsar is very young
2. High-energy positrons lose energy faster than low-energy positrons
3. These initial positrons build up sharp feature in spectrum over time





Spectrum of an Individual Pulsar

1. Large fraction of positrons is produced when pulsar is very young
2. High-energy positrons lose energy faster than low-energy positrons
3. These initial positrons build up sharp feature in spectrum over time




Spectrum of an Individual Pulsar

1. Large fraction of positrons is produced when pulsar is very young
2. High-energy positrons lose energy faster than low-energy positrons
3. These initial positrons build up sharp feature in spectrum over time



Sharp Spectral Features?

- Annihilating dark matter would produce sharp spectral features
- Energy loss processes set up a tension of pulsar feature with dark matter



Energy Loss Rate

Continuous energy loss rate:

$$\frac{dE}{dt} = -\frac{4}{3}\sigma_T \left(\frac{E}{m_e}\right)^2 \left[\rho_B + \sum_i \rho_i(\nu_i) S(E, \nu_i) \right]$$

σ_T : Thomson cross section
 E_e : Electron energy
 m_e : Electron mass
 u_i : ISRF photon energy density
 ν_i : ISRF photon energy
 S : Klein-Nishina suppression

Synchrotron radiation
in magnetic fields

Inverse-Compton scattering
on ambient ISRF photons

Synchrotron Losses

Average energy loss per interaction:

$$E_{\text{crit}} \approx 0.06 \left(\frac{B}{1 \mu\text{G}} \right) \left(\frac{E_e}{1 \text{ TeV}} \right)^2$$

For typical magnetic field strength $B \sim 3 \mu\text{G}$ and electron energy $E_e = 100 \text{ TeV}$


$$\approx 1.8 \text{ keV}$$

Synchrotron losses are small and approximately continuous.

Inverse-Compton Scattering

High energy electrons scatter with photons of the interstellar radiation field

Inverse Compton Scattering



Interstellar Radiation Field (ISRF):

- CMB photons
- IR radiation
- Starlight
- UV radiation

σ_T : Thomson cross section
 E_e : Electron energy
 m_e : Electron mass
 u_i : ISRF photon energy densities
 ν_i : ISRF photon energy


$$\frac{dE_e}{dt} = -\frac{4}{3}\sigma_T c \left(\frac{E_e}{m_e}\right)^2 \sum_i u_i(\nu_i) S_i(E_e, \nu_i)$$

S_i : Klein-Nishina suppression

Inverse-Compton Scattering: Continuous Modelling Fails

Average energy loss per **ICS interaction**

$$\approx \begin{cases} \text{at } E = 1 \text{ TeV} & \rightarrow 0.007 \text{ TeV} \\ \text{at } E = 10 \text{ TeV} & \rightarrow 0.4 \text{ TeV} \\ \text{at } E = 100 \text{ TeV} & \rightarrow 10 \text{ TeV} \end{cases}$$



Modelling Energy Losses

Continuous energy loss rate:

$$\frac{dE}{dt} = -\frac{4}{3}\sigma_T \left(\frac{E}{m_e}\right)^2 \left[\rho_B + \sum_i \rho_i(\nu_i) S(E, \nu_i) \right]$$

Synchrotron radiation
in magnetic fields

Inverse-Compton scattering
on ambient ISRF photons

Approximately continuous.

ICS is a stochastic process
with catastrophic energy losses.

Stochastic Inverse-Compton Scattering Model

[I. John & T. Linden, arXiv:2206.04699]

1. Create positron with some initial energy
2. Evolve in time steps:
 - Calculate **synchrotron** energy losses
 - Based on positron energy, determine if **inverse-Compton scattering** happens and at what photon energy
 - If **ICS**: Calculate energy loss and new positron energy
3. Repeat until desired cooling time is reached


Stochasticity of Inverse-Compton Scattering

Stochastic ICS:

- ICS interactions are rare (~110 interactions in 342 kyr)
- Catastrophic energy losses (~10–100% of energy lost)
- ~30% spread in final positron energy distribution


Continuous calculation:

- All positrons are treated the same way, cool down to exactly the same energy



Positron Spectrum of Individual Pulsar

Example Pulsar:
Geminga
Age: 342 kyr
Distance: 250 pc






Analytic Model:
Hooper et. al,
arXiv:0810.1527



Sharp spectral features introduced by **continuous approximation** are smoothed out by ~50% when correctly treating **inverse-Compton scattering stochastically**

Work in Progress: Spectra For A Range of Pulsar Models

[I. John & T. Linden, arXiv:23xx.xxxx]



Pulsars and Dark Matter Can Be Distinguished




Positron Injection from Pulsars and Dark Matter

Pulsars


Leptophilic Dark Matter

Burst-like injection of e^+e^-



Continuous injection of e^+e^-


Distribution of e^+e^-
injection energies
(power law)



Sharply peaked e^+e^-
injection energy
(corresponding to dark
matter mass)


Catastrophic and Rare Inverse-Compton Scattering

[I. John & T. Linden, arXiv:2304.07317]




ICS interactions are rare, but take a large fraction of the energy in a single interaction, especially at high energies

Energy Loss Times



Energy losses happen slower
in stochastic model than in
continuous model


Enhancement of Dark Matter Signal



Near the dark matter mass, the spectral cutoff is enhanced by about a factor of 2.6


Increased Detectability: Dependence on Energy Resolution

5 % energy resolution




Increased Detectability: Dependence on Energy Resolution


1 % energy resolution
 Expected for e.g. HERD



Dark Matter Signal is Enhanced




5 % energy resolution



1 % energy resolution


Implications of Stochastic ICS

Pulsars do not produce sharp spectral features



[arXiv:2206.04699]

Leptophilic dark matter signal is enhanced




[arXiv:2304.07317]

Dark matter is the only known astrophysical mechanism that can produce sharp spectral features in the e^+e^- flux.

Extra Slides

Dark Matter Annihilation into Muons



- Dark matter annihilates into $\mu^+\mu^-$ that subsequently decay into e^+e^-
- e^+e^- are injected at a distribution of energies
- Enhancement is smaller than in direct e^+e^- case
- Enhancement is further reduced for annihilations into $\tau^+\tau^-$ and other hadronic final states


Stochasticity of Inverse-Compton Scattering

Stochastic ICS:


- ICS interactions are rare (~110 interactions in 342 kyr)
- Catastrophic energy losses (~10–100% of energy lost)
- ~30% spread in final positron energy distribution

Continuous calculation:

- All positrons are treated the same way, cool down to exactly the same energy




Interstellar Radiation Fields and Magnetic Fields



Inverse-Compton Energy Losses



[I. John & T. Linden, arXiv:2304.07317]




ICS interactions are rare, but take a large fraction of the energy in a single interaction, especially at high energies

Implications for Pulsar Models

- Pulsars do not produce sharp features
- Recent papers that fit pulsars to the positron data require large number of pulsars to wash out sharp features below 500 GeV: Possibly only smaller number of pulsars needed to fit positron flux
- Loosens constraints on number of contributing pulsars



Orusa et al., arXiv:2107.06300



Cholis & Krommydas, arXiv:2111.05864