

Dark
matter



F I L E S



Alexander Kusenko (UCLA)

Stockholm, '07

Dark matter's X-files

Dark matter's X-files

- Theory: the seesaw mechanism, *well adjusted*

Dark matter's X-files

- Theory: the seesaw mechanism, *well adjusted*
- Production of sterile neutrinos in the early universe

Dark matter's X-files

- Theory: the seesaw mechanism, *well adjusted*
- Production of sterile neutrinos in the early universe
- Astrophysical hints of sterile dark matter: pulsar velocities, supernova asymmetry, star formation
- Lyman- α bounds on the mass **are** model-dependent
- X-ray bounds and the future prospects: planned Suzaku observations (together with Loewenstein, Biermann)

Sterile neutrinos

The name "sterile" was coined by **Bruno Pontecorvo** in a paper [JETP, **53**, 1717 (1967)], which also discussed

- lepton number violation
- neutrinoless double beta decay
- rare processes (e.g. $\mu \rightarrow e\gamma$)
- vacuum neutrino oscillations
- detection of neutrino oscillations
- astrophysical neutrino oscillations



Бруно Понтекорво



Pontecorvo: neutrino oscillations can "convert potentially active particles into particles that are, from the point of view of ordinary weak interactions, **sterile**, i.e. practically unobservable, since they have the "incorrect" helicity" [JETP, **53**, 1717 (1967)]

Neutrino masses

Discovery of neutrino masses implies a plausible existence of right-handed (sterile) neutrinos. Most models of neutrino masses introduce sterile states

$$\{\nu_e, \nu_\mu, \nu_\tau, \nu_{s,1}, \nu_{s,2}, \dots, \nu_{s,N}\}$$

and consider the following lagrangian:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \bar{\nu}_{s,a} (i\partial_\mu \gamma^\mu) \nu_{s,a} - y_{\alpha a} H \bar{L}_\alpha \nu_{s,a} - \frac{M_{ab}}{2} \bar{\nu}_{s,a}^c \nu_{s,b} + h.c.,$$

where H is the Higgs boson and L_α ($\alpha = e, \mu, \tau$) are the lepton doublets. The mass matrix:

$$M = \begin{pmatrix} \tilde{m}_{3 \times 3} & D_{3 \times N} \\ D_{N \times 3}^T & M_{N \times N} \end{pmatrix}$$

Neutrino masses

Discovery of neutrino masses implies a plausible existence of right-handed (sterile) neutrinos. Most models of neutrino masses introduce sterile states

$$\{\nu_e, \nu_\mu, \nu_\tau, \nu_{s,1}, \nu_{s,2}, \dots, \nu_{s,N}\}$$

and consider the following lagrangian:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \bar{\nu}_{s,a} (i\partial_\mu \gamma^\mu) \nu_{s,a} - y_{\alpha a} H \bar{L}_\alpha \nu_{s,a} - \frac{M_{ab}}{2} \bar{\nu}_{s,a}^c \nu_{s,b} + h.c.,$$

where H is the Higgs boson and L_α ($\alpha = e, \mu, \tau$) are the lepton doublets. The mass matrix:

$$M = \begin{pmatrix} 0 & D_{3 \times N} \\ D_{N \times 3}^T & M_{N \times N} \end{pmatrix}$$

What is the *natural* scale of M ?

Seesaw mechanism

In the Standard Model, the matrix D arises from the Higgs mechanism:

$$D_{ij} = y_{ij} \langle H \rangle$$

Smallness of neutrino masses **does not** imply the smallness of Yukawa couplings. For large M ,

$$m_\nu \sim \frac{y^2 \langle H \rangle^2}{M}$$

One can understand the smallness of neutrino masses even if the Yukawa couplings are $y \sim 1$ [Gell-Mann, Ramond, Slansky; Yanagida; Glashow; Mohapatra, Senjanović].

Is $y \sim 1$ better than $y \ll 1$?

Depends on the model.

- If $y \approx$ some intersection number in string theory, then $y \sim 1$ is natural
- If y comes from wave function overlap of fermions living on different branes in a model with extra-dimensions, then it can be exponentially suppressed, hence, $y \ll 1$ is natural.

In the absence of theory of the Yukawa couplings, one evokes some naturalness arguments.

Clues from cosmology?

Baryon asymmetry of the universe could be generated by **leptogenesis**

However, leptogenesis can work for both $M \gg 100$ GeV and $M < 100$ GeV:

- For $M \gg 100$ GeV, heavy sterile neutrino decays can produce the lepton asymmetry, which is converted to baryon asymmetry by sphalerons [Fukugita, Yanagida]
- For $M < 100$ GeV, neutrino oscillations can produce the lepton asymmetry, which is converted to baryon asymmetry by sphalerons [Akhmedov, Rubakov, Smirnov; Asaka, Shaposhnikov]

Over the years, neutrino physics has shown many theoretical prejudices to be wrong: neutrinos were expected to be massless, neutrinos were expected to have small mixing angles, etc.

Since the fundamental theory of neutrino masses is lacking, one should

**consider all allowed values
for the sterile neutrino masses**

in the following lagrangian:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \bar{\nu}_{s,a} (i\partial_\mu \gamma^\mu) \nu_{s,a} - y_{\alpha a} H \bar{L}_\alpha \nu_{s,a} - \frac{M_{aa}}{2} \bar{\nu}_{s,a}^c \nu_{s,a} + h.c.,$$

where M is can be small or large

Astrophysical clues: dark matter

Dark matter – a simple (minimalist) solution: use one of the particles already introduced to give the neutrino masses

⇒ sterile neutrino

side benefit: explanation of the pulsar kicks, supernova asymmetries

Sterile neutrinos in the early universe

Sterile neutrinos are produced in primordial plasma through

- off-resonance oscillations. [Dodelson, Widrow; Abazajian, Fuller; Dolgov, Hansen; Asaka, Laine, Shaposhnikov et al.]
- oscillations on resonance, if the lepton asymmetry is non-negligible [Fuller, Shi]
- **production mechanisms which do not involve oscillations**
 - inflaton decays directly into sterile neutrinos [Shaposhnikov, Tkachev]
 - Higgs physics: both mass and production [AK]

Active–sterile oscillations

$$\begin{cases} |\nu_1\rangle = \cos\theta|\nu_e\rangle - \sin\theta|\nu_s\rangle \\ |\nu_2\rangle = \sin\theta|\nu_e\rangle + \cos\theta|\nu_s\rangle \end{cases} \quad (1)$$

The almost-sterile neutrino, $|\nu_2\rangle$ was never in equilibrium. Production of ν_2 could take place through oscillations.

The coupling of ν_2 to weak currents is also suppressed, and $\sigma \propto \sin^2\theta$.

The probability of $\nu_e \rightarrow \nu_s$ conversion in presence of matter is

$$\langle P_m \rangle = \frac{1}{2} \left[1 + \left(\frac{\lambda_{\text{osc}}}{2\lambda_s} \right)^2 \right]^{-1} \sin^2 2\theta_m, \quad (2)$$

where λ_{osc} is the oscillation length, and λ_s is the scattering length.

Mixing is suppressed at high temperature [Dolgov, Barbieri; Kainulainen; Stodolsky]

$$\sin^2 2\theta_m = \frac{(\Delta m^2/2p)^2 \sin^2 2\theta}{(\Delta m^2/2p)^2 \sin^2 2\theta + (\Delta m^2/2p \cos 2\theta - V(T))^2}, \quad (3)$$

For small angles,

$$\sin 2\theta_m \approx \frac{\sin 2\theta}{1 + 0.79 \times 10^{-13} (T/\text{MeV})^6 (\text{keV}^2/\Delta m^2)} \quad (4)$$

Production of sterile neutrinos peaks at temperature


$$T_{\max} = 130 \text{ MeV} \left(\frac{\Delta m^2}{\text{keV}^2} \right)^{1/6}$$

The resulting density of relic sterile neutrinos in conventional cosmology, in the absence of a large lepton asymmetry:

$$\Omega_{\nu_2} \sim 0.3 \left(\frac{\sin^2 2\theta}{10^{-8}} \right) \left(\frac{m_s}{\text{keV}} \right)^2$$

[Dodelson, Widrow; Abazajian, Fuller, Patel; Dolgov, Hansen; Fuller, Shi]

Hadronic uncertainties under control [Asaka, Laine, Shaposhnikov]




Lepton asymmetry and the MSW resonance

If the lepton asymmetry L is non-zero, sterile neutrinos can be produced on resonance.


[Fuller, Shi; Abazajian, Fuller, Patel]

The amount of dark matter and the momentum distribution depend on L .






Lepton asymmetry and the MSW resonance

MSW resonance favors the low-momentum neutrinos. The dotted line is the normalized active neutrino spectrum. The thick-solid, dashed, dot-dashed lines correspond to $L = 0.01$, mass around 1 keV, and different mixing angles. [Fuller, Shi; Abazajian, Fuller, Patel]



Dark matter and the Lyman- α forest.



Dark matter and the Lyman- α forest.


The bounds depend on the production mechanism.

$$\lambda_{FS} \approx 1 \text{ Mpc} \left(\frac{\text{keV}}{m_s} \right) \left(\frac{\langle p_s \rangle}{3.15 T} \right)_{T \approx 1 \text{ keV}}$$

The ratio

$$\left(\frac{\langle p_s \rangle}{3.15 T} \right)_{T \approx 1 \text{ keV}} = \begin{cases} 0.9 & \text{for production off - resonance} \\ 0.6 & \text{for MSW resonance (depends on } L) \\ 0.2 & \text{for production at } T > 100 \text{ GeV} \end{cases}$$

For DW production, is Ly- α in conflict with dSphs?



[Viel et al.; Seljak et al., Gilmore et al.]

Neutrino masses: new scale or new Higgs physics?

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \bar{N}_a (i\partial_\mu \gamma^\mu) N_a - y_{\alpha a} H \bar{L}_\alpha N_a - \frac{M_a}{2} \bar{N}_a^c N_a + h.c.,$$

To explain the pulsar kicks and dark matter, one needs $M \sim \text{keV}$. Is this a new fundamental scale? Perhaps. Alternatively, it could arise from the Higgs mechanism:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \bar{N}_a (i\partial_\mu \gamma^\mu) N_a - y_{\alpha a} H \bar{L}_\alpha N_a - h_a S \bar{N}_a^c N_a + V(H, S)$$

$$M = h \langle S \rangle$$

Now $S \rightarrow NN$ decays can produce sterile neutrinos

For small h , the sterile neutrinos are out of equilibrium in the early universe, but S is in equilibrium. There is a new mechanism to produce sterile dark matter at $T \sim m_S$ from decays $S \rightarrow NN$:


$$\Omega_s = 0.2 \left(\frac{33}{\xi} \right) \left(\frac{h}{1.4 \times 10^{-8}} \right)^3 \left(\frac{\langle S \rangle}{\tilde{m}_s} \right)$$

Here ζ is the dilution factor due to the change in effective numbers of degrees of freedom.

$$\langle S \rangle = \frac{M_s}{h} \sim \frac{\text{few keV}}{1.4 \times 10^{-8}} \sim 10^2 \text{ GeV}$$

The sterile neutrino momenta are red-shifted by factor $\zeta^{1/3} \approx 3$.

Cooling changes the bounds



[AK, PRL **97**:241301 (2006)]

Implications for the EW phase transition and the LHC

The presence of S in the Higgs sector changes the nature of the electroweak phase transition, which now proceeds in two stages:

$$\{S = 0, H = 0\} \longrightarrow \{S \neq 0, H = 0\} \longrightarrow \{S \neq 0, H \neq 0\}$$

One may be able to discover the *invisible Higgs* at the LHC in the $Z + H_{\text{inv}}$ channel, as well as in the weak boson fusion channel. In some range of masses, the discovery is possible at the LHC with 10 fb^{-1} in the $Z + H_{\text{inv}}$ channel [Davoudiasl et al.] LHC phenomenology [O'Connell et al.]

Astrophysical clues: supernova

- Sterile neutrino emission from a supernova is anisotropic due to
 1. asymmetries in the urca cross sections
 2. magnetic effects on neutrino oscillations
- Sterile neutrinos with masses and mixing angles consistent with dark matter can explain the pulsar velocities

[AK, Segrè; Fuller, AK, Mocioiu, Pascoli; Barkovich, D'Olivo, Montemayor]

The pulsar velocities.

Pulsars have large velocities, $\langle v \rangle \approx 250 - 450 \text{ km/s}$.

[Cordes *et al.*; Hansen, Phinney; Kulkarni *et al.*; Lyne *et al.*]

A significant population with $v > 700 \text{ km/s}$,


about **15 %** have $v > 1000 \text{ km/s}$, up to **1600 km/s**.

[Arzoumanian *et al.*; Thorsett *et al.*]

A very fast pulsar in Guitar Nebula




HST, December 1994



HST, December 2001



Map of pulsar velocities



Proposed explanations:

- asymmetric collapse [Shklovskii] (small kick)
- evolution of close binaries [Gott, Gunn, Ostriker] (not enough)
- acceleration by EM radiation [Harrison, Tademaru] (kick small, predicted polarization not observed)
- asymmetry in EW processes that produce neutrinos [Chugai; Dorofeev, Rodinov, Ternov] (asymmetry washed out)
- “cumulative” parity violation [Lai, Qian; Janka] (it’s *not* cumulative)
- various exotic explanations
- explanations that were “not even wrong” ...

Asymmetric collapse



“...the most extreme asymmetric collapses do not produce final neutron star velocities above 200km/s” [Fryer '03]

Supernova neutrinos

Nuclear reactions in stars lead to a formation of a heavy iron core. When it reaches $M \approx 1.4M_{\odot}$, the pressure can no longer support gravity. \Rightarrow collapse.

Energy released:

$$\Delta E \sim \frac{G_N M_{\text{Fe core}}^2}{R} \sim 10^{53} \text{erg}$$

99% of this energy is emitted in neutrinos

Pulsar kicks from neutrino emission?

Pulsar with $v \sim 500$ km/s has momentum

$$M_{\odot} v \sim 10^{41} \text{ g cm/s}$$

SN energy released: 10^{53} erg \Rightarrow in neutrinos. Thus, the total neutrino momentum is

$$P_{\nu; \text{total}} \sim 10^{43} \text{ g cm/s}$$

a **1% asymmetry** in the distribution of **neutrinos**

is sufficient to explain the pulsar kick velocities

But what can cause the asymmetry??

Magnetic field?

Neutron stars have large magnetic fields. A typical pulsar has surface magnetic field $B \sim 10^{12} - 10^{13} \text{ G}$.

Recent discovery of *soft gamma repeaters* and their identification as *magnetars*


⇒ some neutron stars have surface magnetic fields as high as $10^{15} - 10^{16} \text{ G}$.

⇒ magnetic fields inside can be $10^{15} - 10^{16} \text{ G}$.

Neutrino magnetic moments are negligible, but the **scattering of neutrinos off polarized electrons and nucleons** is affected by the magnetic field.


Core collapse supernova

Onset of the collapse: $t = 0$



Core collapse supernova


Shock formation and “neutronization burst”: $t = 1 - 10$ ms



Protoneutron star formed. Neutrinos are trapped. The shock wave breaks up nuclei, and the initial neutrino come out (a few %).

Core collapse supernova

Thermal cooling: $t = 10 - 15$ s



Most of the neutrinos emitted during the cooling stage.

Electroweak processes producing neutrinos (urca),

$$p + e^- \rightleftharpoons n + \nu_e \text{ and } n + e^+ \rightleftharpoons p + \bar{\nu}_e$$

have an asymmetry in the production cross section, depending on the spin orientation.


$$\sigma(\uparrow e^-, \uparrow \nu) \neq \sigma(\uparrow e^-, \downarrow \nu)$$

The asymmetry:

$$\tilde{\epsilon} = \frac{g_V^2 - g_A^2}{g_V^2 + 3g_A^2} k_0 \approx 0.4 k_0,$$

where k_0 is the fraction of electrons in the lowest Landau level.

In a strong magnetic field,




k_0 is the fraction of electrons in the lowest Landau level.

Pulsar kicks from the asymmetric production of neutrinos?

[Chugai; Dorofeev, Rodionov, Ternov]

Can the weak interactions asymmetry cause an anisotropy in the flux of neutrinos due to a large magnetic field?



Neutrinos are trapped at high density.

Can the weak interactions asymmetry cause an anisotropy in the flux of neutrinos due to a large magnetic field?


No

Rescattering washes out the asymmetry


In approximate thermal equilibrium the asymmetries in scattering amplitudes do not lead to an anisotropic emission [Vilenkin,AK, Segrè]. Only the outer regions, near neutrinospheres, contribute, but the kick would require a mass difference of $\sim 10^2$ eV [AK,Segrè].

However, if a weaker-interacting sterile neutrino was produced in these processes, the asymmetry would, indeed, result in a pulsar kick!

[AK, Segrè; Fuller, AK, Mocioiu, Pascoli]



Allowed range of parameters (time scales, fraction of total energy emitted):



[Fuller, AK, Mocioiu, Pascoli]

Resonance in the magnetic field

Matter potential:

$$V(\nu_s) = 0$$


$$V(\nu_e) = -V(\bar{\nu}_e) = V_0 (3Y_e - 1 + 4Y_{\nu_e})$$

$$V(\nu_{\mu,\tau}) = -V(\bar{\nu}_{\mu,\tau}) = V_0 (Y_e - 1 + 2Y_{\nu_e}) + c_L^z \frac{\vec{k} \cdot \vec{B}}{k}$$

$$c_L^z = \frac{eG_F}{\sqrt{2}} \left(\frac{3N_e}{\pi^4} \right)^{1/3}$$


[D'Olivo, Nieves, Pal; Semikoz]

The magnetic field shifts the position of the resonance because of the $\frac{\vec{k} \cdot \vec{B}}{k}$ term in the potential:



In the absence of magnetic field, ν_s escape isotropically


The magnetic field shifts the position of the resonance because of the $\frac{\vec{k} \cdot \vec{B}}{k}$ term in the potential:



Down going neutrinos have higher energies


of the

The range of parameters for off-resonance transitions:




[AK, Segrè]

Resonance & off-resonance oscillations




[A.K., Segrè; Fuller, A.K., Mocioiu, Pascoli; Barkovich, D'Ollivo, Montemayor]

Allowed range of masses and mixng angles




Other predictions of the pulsar kick mechanism

- Stronger supernova shock [Fryer, AK]




Other predictions of the pulsar kick mechanism

- Stronger supernova shock [Fryer, AK]




Other predictions of the pulsar kick mechanism

- Stronger supernova shock [Fryer, AK]
- **No $B - v$ correlation** expected because
 - the magnetic field *inside* a hot neutron star during the *first ten seconds* is very different from the surface magnetic field of a cold pulsar
 - rotation washes out the x, y components
- **Directional $\vec{\Omega} - \vec{v}$ correlation** is expected, because
 - the direction of rotation remains unchanged
 - only the z -component survives




Radiative decay

Sterile neutrino in the mass range of interest have lifetimes **longer than the age of the universe**, but they do decay:




Photons have energies $m/2$: X-rays. Large lumps of dark matter emit some X-rays. [Abazajian, Fuller, Tucker; Dolgov, Hansen; Shaposhnikov et al.]



X-ray observations: the current limits



X-ray observations: Draco and Ursa Minor




X-ray observations: Suzaku reach



[Loewenstein, Biermann, AK]

Astrophysical clues: star formation and reionization

Molecular hydrogen is necessary for star formation




[Tegmark, et al., ApJ 474, 1 (1997)]

Molecular hydrogen




In the presence of ions the following reactions are faster:



H^+ catalyze the formation of molecular hydrogen


[Biermann, AK, PRL **96**, 091301 (2006)]

[Stasielak, Biermann, AK, ApJ.654:290 (2007)]



[Biermann, AK; Stasielak, Biermann, AK]

Clues of sterile neutrinos



*This could be the greatest discovery of the century.
Depending, of course, on how far down it goes.*

Summary

- Sterile neutrinos almost certainly exists and have masses between eV and the Planck scale.
- A rather minimal extension of the Standard Model, the addition of **sterile neutrinos**, explains all the present data, including
 - dark matter (warm or cold, depending on the mass)
 - baryon asymmetry of the universe
 - pulsar velocities
- X-ray telescopes (perhaps, Suzaku) can explore the entire region of concordance in the parameter space

