

Effects of topological heterogeneity on random matrix spectra

Peter Sollich

Diego Tapias, Fernando Metz, Jeferson Da Silva

Phys. Rev. Res. 2: 043116, 2020

J. Phys. Complexity 3: 045012, 2022

Europhys. Lett. 144: 41001, 2023

SciPost Physics 18: 047, 2025 + unpublished


Glassy systems



Disordered solids: dense & amorphous

Making a glass


Monodisperse Liquid




Temperature
Reduction



Crystal




Polydisperse Liquid




Temperature
Reduction



Glass



Configuration space dynamics



Doye 2002

- Rough energy landscape, many minima
- Low temperature dynamics: activated hops
- **Effective network** of states $i = 1, \dots, N$

Master operator


- **Probability** of being in state i : $p_i(t)$
- **Hopping rate** $i \rightarrow j$: W_{ji}
- **Network structure**: $a_{ij} = 1$ if edge present, else $= 0$
- Time evol'n: $\frac{d}{dt}p_i = \sum_j a_{ij}(W_{ij}p_j - W_{ji}p_i) = - \sum_j L_{ij}p_j$
- Master operator L is **graph Laplacian** $L = D - A$
- $A_{ij} = a_{ij}W_{ij}$: off-diagonal, (weighted) **adjacency matrix**
- $D_{ii} = \sum_j W_{ji}$: balances diagonal terms

Quantum analogue


- Corresponding quantum Hamiltonian:

$$H = \sum_i \epsilon_i c_i^\dagger c_i + \sum_{(ij)} W_{ij} (c_i^\dagger c_j + \text{h.c.}), \quad \epsilon_i = -D_{ii}$$

- Similar to Anderson model on network, widely studied e.g. for **Many-Body Localisation**
- Disorder** also in hopping terms, **correlated** with on-site terms
- For now, ignore correlations



Network model & disorder




Network model & disorder

- **Focus:** Effects of **network topology**
Classically: e.g. from kinetic constraints in real space
Quantum: Fock space connectivity
- Network model: fix **distribution** p_k of **degrees** $k_i = \sum_j a_{ij}$
- Network otherwise random: configuration model
- **Connectivity** $c = \langle k \rangle =$ average degree
- Weights: Gaussian with zero mean and $\langle W_{ij}^2 \rangle = 1/c$

Topological heterogeneity

- Study limit of **high connectivity** $c = \langle k \rangle$
Why? Physical glass in volume V has $N \sim e^V$, $c \sim V \sim \ln N$
- Degree distribution p_k ? For Erdős-Renyí graphs, Poisson distribution, relative variance $c/c^2 \rightarrow 0$ for $c \rightarrow \infty$
- Heterogeneity effects disappear for $c \rightarrow \infty$, effectively random regular graph


(Cugliandolo Schehr Tarzia
Venturelli 2024)



Topological heterogeneity

- Alternative choices for degree distribution p_k ?
- Negative binomial distribution:
scaled degrees $\kappa = k/c$ have Gamma-distribution $\nu(\kappa) \propto \kappa^{\gamma-1} e^{-\gamma\kappa}$
- Variance $1/\gamma$ (relative to mean $\langle \kappa \rangle = 1$) does not vanish for $c \rightarrow \infty$
- Small $\gamma \Leftrightarrow$ heterogeneous networks
- Study spectral density $\rho(E) = \langle \text{Im } G_{ii} \rangle / \pi$ of eigenvalues E of H
- Local density of states $\text{Im } G_{ii}$ from resolvent $G = (z - H)^{-1}$,
with $z = E - i\eta$

Cavity method







- Network becomes **locally tree-like** for $N \rightarrow \infty$, can use **cavity method** to get recursive equations for cavity resolvent entries $G_{ii}^{(j)}$
- Large c : self-consistent equation for average of $G_{ii}^{(j)}$
- Limits: $\lim_{\eta \rightarrow 0} \lim_{c \rightarrow \infty} \lim_{N \rightarrow \infty} \dots$





Part 1: No on-site disorder

$$H = \sum_{(ij)} W_{ij} (c_i^\dagger c_j + \text{h.c.})$$

Spectral density







Spectral density



Wigner


Spectral density

Spectral
divergence
 $\rho(E) \sim |E|^{\gamma-1}$



Wigner


Free multiplication



- Can show: spectral density = **free multiplicative convolution** (Dembo+ 2021)
- Effective decomposition: $H = D^{1/2} W D^{1/2}$, $D_{ii} = \kappa_i$

Local density of states


Distribution of $y_i = \text{Im } G_{ii}$ at $E \neq 0$



- Distribution of y has **finite support**
- Eigenvectors are extended

Local density of states

Distribution of $y_i = \text{Im } G_{ii}$ at $E = 0$



Multifractality

- Characterise structure of eigenvectors
- Generalised **inverse participation ratio** (IPR):

$$I_q(N) = \sum_i |\psi_i|^{2q} \sim N^{-\tau(q)}$$


- **Multifractal exponents** $\tau(q)$:
Extended state $|\psi_i|^2 \sim 1/N$ gives $\tau(q) = q - 1$
Localised state $|\psi_i|^2 = O(1)$ (for $O(1)$ nodes): $\tau(q) = 0$
Single fractal dimension: $\tau(q) = d_f(q - 1)$
- Evaluate I_q from scaling of LDoS distribution
("envelope distribution", De Luca+ 2014)

Multifractal exponents

$$E = 0, \gamma < 1$$

Eigenvectors
localised with
multifractal
features:

$$\tau = \begin{cases} q/\gamma - 1 \\ 0 \end{cases}$$




Multifractal exponents

$$E = 0, \gamma > 1$$

Eigenvectors
extended non-
ergodic:


$$\tau = \begin{cases} q - 1 \\ q(1 - 1/\gamma) \end{cases}$$

(for $q \lesssim \gamma$)



Wavefunction amplitudes

- Local density of states is
$$y_i \propto \sum_{\mu} |\psi_i^{\mu}|^2 \delta_{\eta}(E - E_{\mu})$$
- Expect similar features in distribution of **squared wavefunction amplitudes**
$$x_i = N |\psi_i|^2$$
(see e.g. Monthus 2017)




Statistical localisation

- What is **mechanism of localisation** at $E = 0$?
- Not power law in “space” on network, or competing correlation length scales
(Kravtsov+ 2015, Monthus 2016; García-Mata+ 2020, 2022, Das+ 2025, ...)
- Strong correlation between $|\psi_i|^2$ and **degree** k_i
- Localisation determined by **statistical** properties of nodes

Statistical localisation

- What
- Not
corr
(Kraus)
- Struc
- Loc




Part 2:

Adding on-site disorder

$$H = \sum_i \epsilon_i c_i^\dagger c_i + \sum_{(ij)} W_{ij} (c_i^\dagger c_j + \text{h.c.}), \quad |\epsilon_i| \leq \frac{1}{2}W$$


What changes?

- Spectral density $\rho(E)$: **no divergences**, still has unbounded support
- LDoS distribution now has large- y power law tail
 $P(y) \sim y^{-\gamma-2}$
for **all** $|E| < \frac{1}{2}W$
- For $|E| > W/2$, LDoS has compact support again




What changes?

- LDoS distribution acquires additional **small- y power law tail** $P(y) \sim \nu(y) \sim y^{\gamma-1}$ (for all E)




Wavefunction amplitudes




Spectrum of fractal dimensions

$P(y) \sim \nu(y)$ vs $P(y) \sim y^{-3}\nu(1/y)$: Mirlin-Fyodorov **symmetry**




$$f(\alpha) = \begin{cases} -\gamma + (1+\gamma)\alpha & \frac{\gamma}{1+\gamma} \leq \alpha \leq 1 \\ \gamma + 1 - \gamma\alpha & 1 \leq \alpha \leq \frac{\gamma+1}{\gamma} \end{cases}$$



$$\tau_q = \begin{cases} q - 1 & q < 1 + \gamma \\ \left(\frac{\gamma}{1+\gamma}\right)q & q > 1 + \gamma \end{cases}$$

Phase diagram



Part 2b:

Power-law on-site disorder


$$p_{\epsilon}(\epsilon) \sim |\epsilon|^{\beta-1}$$

What changes?

LDoS distribution
acquires distinct
large- y power-law
tail for $E = 0$:

$$P(y) \sim \frac{\nu(1/y)}{y^3} p_\epsilon(1/y)$$

$$P(y) \sim y^{-(\beta+\gamma+1)}$$




Part 3:

Graph Laplacians


$$H = \sum_i \epsilon_i c_i^\dagger c_i + \sum_{(ij)} W_{ij} (c_i^\dagger c_j + \text{h.c.})$$

$$\epsilon_i = -D_{ii} = -\sum_j W_{ji}$$

Spectral density




Spectral density



Local density of states


Distribution of $y_i = \text{Im } G_{ii}$ at $E \neq 0$

- Distribution of y has **exponential tail**
- Eigenvectors are extended




Local density of states

Distribution of $y_i = \text{Im } G_{ii}$ at $E = 0$




Local density of states

Distribution of $y_i = \text{Im } G_{ii}$ at $E = 0$



Wavefunction amplitudes

Distribution of $|\psi_i|^2$ at $\lambda = 0$




Multifractal exponents

$$E = 0, \gamma < 1/2$$

Eigenvectors
localised with
multifractal features:

$$\tau = \begin{cases} \frac{q}{\gamma + 1/2} - 1 \\ 0 \end{cases}$$



For $\gamma > 1/2$, extended non-ergodic: $\tau = \begin{cases} q - 1 \\ q(\gamma - 1/2)/(\gamma + 1/2) \end{cases}$

Summary & Outlook

- **Heterogeneous** networks can “break” Wigner law even for $c \rightarrow \infty$
- **Localisation transitions**, **multifractal** eigenvectors, with on-site disorder for $|E| < W/2$, else at spectral divergence
- Nontrivial differences between A and L ($\gamma \rightarrow \gamma + \frac{1}{2}$), connections to classical trajectory biasing?
- **Statistical** rather than spatial localisation, can occur in other models on deep minima
- **Origin** of M-F **symmetry**?
- **“Strong”** multifractality?

